

Q B

741


E2

UC-NRLF



\$B 560 383

IN MEMORIAM
George Davidson
1825-1911



EX LIBRIS

Compliments of the a

Recd July 28/90

Comp. Dan. Jr

PHILOSOPHICAL SOCIETY OF WASHINGTON

BULLETIN VOL. XI, pp. 275-358.

THE PROGRESS

OF

METEORIC ASTRONOMY

IN

AMERICA.

BY

JOHN ROBIE EASTMAN.

READ BEFORE THE PHILOSOPHICAL SOCIETY OF WASHINGTON,
APRIL 12, 1890.

WASHINGTON:
PUBLISHED BY THE SOCIETY
JULY, 1890.

Davidson bookplate

THE PROGRESS OF METEORIC ASTRONOMY IN AMERICA.

BY

JOHN ROBIE EASTMAN.

[Read before the Society April 12, 1890.]

TABLE OF CONTENTS.

	Page.
Introduction.....	276
Abstracts of Theories.....	278
Authors:	
Rev. Thomas Clap.....	278
Dr. W. G. Reynolds.....	278
Prof. Edward Hitchcock.....	279
Prof. D. Olmstead.....	279
E. C. Herriek.....	281
Prof. Benjamin Peirce.....	281
Prof. S. C. Walker.....	281
Prof. Peter A. Browne.....	282
Prof. J. Lawrence Smith.....	283
Dr. B. A. Gould.....	283
Prof. H. A. Newton.....	285
Prof. A. C. Twining.....	285
Prof. H. A. Newton.....	285
Prof. A. C. Twining.....	286
B. V. Marsh.....	286
Prof. H. A. Newton.....	287
Prof. Daniel Kirkwood.....	290
Prof. H. A. Newton.....	290
Prof. S. Newcomb.....	291
Prof. W. Harkness.....	291
Prof. Daniel Kirkwood.....	291
Jacob Ennis.....	291
Prof. Pliny Earle Chase.....	292
Prof. H. A. Newton.....	292
Prof. J. W. Mallet.....	292

	Page.
Authors :	
Prof. A. W. Wright	292
Prof. H. A. Newton	292
Prof. H. A. Newton	297
Examination of Theories	298
Meteors	298
Comets	300
Comets and Meteors	302
Lockyer's Theories	305
Huggins on the Spectra of the Aurora and of Nebulæ	310
Liveing and Dewar on the Spectra of Nebulæ and of Magnesium	310
Conclusions	311
Observations of Meteors	312
Catalogues	313
I. Observed Meteorites	316
II. Discovered Meteorites, date of discovery given	318
III. Discovery of Meteorites, date of discovery unknown	322
IV. Meteor Showers	324
V. Sporadic Meteors	336

INTRODUCTION.

The progress of Meteoric Astronomy through its successive stages of development has been so peculiar in America, especially in the United States, that, unlike almost all the other branches of Astronomy and Physics, its advance may be thoroughly discussed with very little reference to the important growth which it has made in Europe.

From the nature of the phenomena it is evident that the apparition and fall of meteors must have compelled the attention of mankind through all ages, but the earliest records are at least obscure. While there may be some claim to authenticity in the early allusions to what was apparently meteoric phenomena, there seem to be no trustworthy observations until about 600 B. C.

From that time the falls of a great number of meteors and meteorites were recorded with more or less accuracy and detail, but no special attention was attracted to the observation and study of such phenomena until the publication of a paper in 1794, by Chladni, on a mass of meteoric iron

found in Siberia by Dr. Pallas, a well-known naturalist. About this time several noted meteorites fell in Europe, and in 1802 Edward Howard published in the *Philosophical Transactions* a paper entitled "Experiments and observations on certain stony substances which at different times are said to have fallen on the earth." This paper contains, probably, the account of the first chemical analysis of a meteorite ever made.

Nearly all the publications referring to meteors, both in Europe and America, up to the year 1833 were confined to vague theories and brief speculations with regard to their origin.

The very important meteoric shower on the morning of April 20, 1803, was the first well-defined phenomenon of that class in this country of which there seems to be any record. There is no evidence that it was well observed except at Portsmouth, N. H., and at Richmond, Va., and no recurrence of this shower of any notable magnitude has since been observed. Graphic accounts of this phenomenon were printed in "The New Hampshire Gazette," of Portsmouth, N. H., May 31, 1803, and in "The Virginia Gazette and General Advertiser," of Richmond, Va., May 23, 1803, but apparently no scientific interest or discussion was developed.

The wonderful display of meteors on the morning of November 14, 1833, which was seen throughout the Atlantic coast of the United States, gave a decided impulse to the study of the subject and suddenly brought the principal American observers into prominence.

The serious study of meteoric phenomena in America may be said to date from this epoch.

The earliest studies immediately developed theories, more or less fantastic, to account for the varied but startling display in the heavens.

The first theories, derived from only a few facts, naturally presented the greatest range of speculation.

As phenomena multiplied, the limits of speculation were

notably contracted, and in 1834 the germ of the true theory of meteoric motion was presented, but not developed.

The most accurate idea of the progress of the science of Meteoric Astronomy can be obtained, without doubt, from an examination of the principal theories.

ABSTRACTS OF THEORIES.

The following abstracts of these theories are presented in chronological order, and in each case the language of the author is employed if practicable.

Probably the first paper printed in this country which advances any theory of the nature or the motion of meteors was written by *Rev. Thomas Clap*, ex-president of Yale College, and was printed in Norwich, Connecticut, in 1781. He concluded that "our observations have heretofore been so imperfect as that we cannot easily determine minute circumstances; but the general theory seems highly probable, if not certain, that these superior meteors are solid bodies, half a mile in diameter, revolving around the earth in long ellipses, their least distance being about twenty or thirty miles; that by their friction upon the atmosphere they make a constant rumbling noise and collect electrical fire, and when they come nearest the earth or a little after, being then overcharged, they make an explosion as loud as a large cannon."

In 1819 *W. G. Reynolds, M. D.*, of Middletown Point, N. J., published a paper¹ advocating the theory that "Meteors proceed from the earth. They arise from certain combinations of its elements with solar heat, and meteoric stones are the necessary results of the decomposition of these combinations."

After the shower of 1833 elaborate accounts of the event were written by several scientific observers, and various conclusions and theories were deduced.

Prof. Edward Hitchcock,² of Amherst College, Mass., con-

¹ A. J. S., I, 266.

² A. J. S., XXV, 354.

cluded that "there was a point from which most of the meteors seemed to emanate ; that this radiant corresponded to that point in the dome of the heavens to which the magnetic needle would point if left free to move vertically and horizontally, and that meteors are only modifications of the Aurora Borealis."

Prof. D. Olmstead,¹ of Yale College, discussed at length the meteors of November 13, 1833, with the following conclusions :

"1st. The meteors originated beyond the limits of our atmosphere and fell towards the earth, in straight and nearly parallel lines, from a point 2,238 miles above the surface of the earth.

"2d. Their velocity on entering the earth's atmosphere was about four miles per second:

"3d. They consisted of light, transparent, combustible matter, and took fire and were consumed in traversing the atmosphere."

Prof. Olmstead finally concluded that "the meteors of November 13 consisted of portions of the extreme parts of a nebulous body which revolves around the sun in an orbit interior to that of the earth, but little inclined to the plane of the ecliptic, having its aphelion near to the earth's path and having a periodic time of 182 days, nearly."

After discussing the November meteors of 1836, Prof. Olmstead² concluded that "the zodiacal light might be the source of those meteors, and therefore was not a portion of the sun's atmosphere, but a nebulous or cometary body revolving around the sun within the earth's orbit nearly in the plane of the solar equator, approaching at times very near to the earth, and having a periodic time of either one year or half a year, nearly."

On the 28th of April, 1840, *Mr. E. C. Herrick*³ read before the Connecticut Academy of Arts and Sciences a paper on

¹ A. J. S., XXVI., 132.

² A. J. S., XXXI., 386.

³ A. J. S., XL., 349.

"The history of star-showers of former times," in which he presented a brief account of all the records he had been able to find, together with the following tabular chronological account of star-showers, where the dates are reduced to Gregorian style:

Chronological List of Star-Showers.

Number.	Date.	Number.	Date.
1	B. C. 1768.	21	A. D. 1060.
2	B. C. 686.	22	" 1090.
3	A. D. 7.	23	" 1094.
4	" 532.	24	" 1095, April 10.
5	" 558.	25	" 1096, April 10 (?).
6	" 585, September 6 (?).	26	" 1106, February 19.
7	" 611.	27	" 1122, April 11.
8	" 744 or 747.	28	" 1199, October (?).
9	" 750.	29	" 1202, October 26.
10	" 764 (?), March.	30	" 1243, August 2.
11	" 765, January 8.	31	" 1366, October 30.
12	" 829.	32	" 1398.
13	" 855, October 21.	33	" 1399, October (?).
14	" 899, November 18.	34	" 1635 or 1636.
15	" 901, November 30.	35	" 1743, October 15.
16	" 902, October 30.	36	" 1799, November 12.
17	" 912 or 913.	37	" 1803, April 20.
18	" 931 or 934, October 19.	38	" 1832, November 13.
19	" 935, October (?).	39	" 1833, November 13.
20	" 1029, July or August.		

Of the theory of meteors, Mr. Herrick wrote :

"The most probable hypothesis is that there are revolving around the sun millions of small planetary and nebulous bodies of various magnitudes and densities, and that when any of these dart through our atmosphere they become ignited and are seen as shooting-stars."

In discussing a paper on meteors by Prof. Erman (*Schumacher's Ast. Nach.* No. 385) *Prof. Benjamin Peirce*,¹ after pointing out an error in Erman's work, concludes in these words : "The plane of the meteors cannot differ much from that of the ecliptic, and their relative velocity cannot exceed one-third of the earth's velocity. A ring so nearly in the plane of the earth's orbit must be subject to great perturbations ; and, if there is one, I think that no observations which we can make will enable us to calculate its motions with any degree of accuracy."

On January 15, 1841, *Prof. S. C. Walker*² read a paper before the American Philosophical Society, "On the periodical meteors of August and November," in which the following points were discussed :

The relative velocities of meteors ;

The relative directions of meteors in space ;

The periodical or anniversary display of meteors ;

The respective plausibilities of the hypotheses of a single cluster with a half-yearly or yearly period, or that of a continuous ring for the periodical meteors of August and November ;

The theories of aerolites and shooting-stars ;

The variation of the relative velocity and of the convergent point ;

And principally the investigation of formulæ for computing the elliptic elements of the orbit of a meteor from its observed relative velocity and direction.

In 1844 an "Essay on Solid Meteors and Meteoric Stones"

¹ Trans. Am. Phil. Soc., VIII₂, 83.

² Trans. Am. Phil. Soc., VIII₂, 87.

was published by *Prof. Peter A. Browne*, of La Fayette College.

The author devoted the first and larger portion of his paper to proving the solidity of meteors.

The latter portion of his essay was confined to the examination and rejection of all the theories previously advanced, which, briefly stated, were :

1st. Dr. Halley's theory that meteors were nothing but a stratum of inflammable vapor, gradually raised from the earth and accumulated in an elevated region, which suddenly took fire at one end and the progress of the flame along the stratum produced the apparent motion of the meteor.

2d. The theory in Luke Howard's Meteorology that hydrogen gas dissolves various bodies, even iron, and that is evolved, mixed with carbon in the gaseous state, from the earth in large quantities, is collected in vast fields in the air, is fired by electric explosions, and, the gasses burning out, they let fall the earthy and metallic contents precipitated and agglutinated as we find them in aerolites.

3d. Prof. Soldani's theory that meteoric stones are generated in the air by a combination of mineral substances which had risen as exhalations from the earth.

4th. Dr. Reynolds' theory, previously given in this paper.

5th. Dr. Blagden's theory that meteors are electrical phenomena.

6th. The theory of Patrick Murray that meteors originate in the local atmosphere of the earth, and their explosions are due to electrical action.

7th. The theories of Brewster and La Grange that meteors are bodies thrown off from the earth by volcanic action.

8th. The theories of Hutton and La Place that meteors are thrown from volcanoes in the moon.

9th. Newton's theory that they proceed from the tail of a comet.

10th. They are terrestrial comets—a theory maintained by Professors Clap and Day and by Carvallo.

11th. The theory that they were solids that have been floating in space from the beginning; advocated by Chladni, Franklin, and Rittenhouse.

12th. The theory of Olbers that they are fragments of an exploded planet.

13th. The theory of Quetelet that they belong to a zone through which the earth passes annually.

14th. The theory of Boubée that they are fragments of an exploded comet.

These theories are all rejected as disproved or absurd; but the author advances no theory as a substitute. He announces, however, that, to his mind, "the most probable supposition yet made is that the solid meteors may possibly emanate from the sun," though no serious attempt is made to prove the proposition.

In a paper read by *Prof. J. Lawrence Smith* before the American Association for the Advancement of Science, in April, 1854, the author¹ advocated the theory of the lunar origin of meteors, which he stated as follows: "The moon is the only large body in space, of which we have any knowledge, possessing the requisite conditions demanded by the physical and chemical properties of meteorites; and they have been thrown off from that body by volcanic action (doubtless long since extinct), and, encountering no gaseous medium of resistance, reached such a distance as that the moon exercised no longer a preponderating attraction, the detached fragment possessing an orbital motion and an orbital velocity which it had in common with all parts of the moon, but now more or less modified by the projectile force and new condition of attraction in which it was placed in reference to the earth, acquired an independent orbit more or less elliptical. This orbit, necessarily subject to great disturbing influences, may sooner or later cross our atmosphere and be intercepted by the body of the globe."

In 1859² *Dr. B. A. Gould* read a paper before the Ameri-

¹ A. J. S., XIX₂, 343.

² Proc. A. A. A. S. 1859, 181.

can Association for the Advancement of Science to disprove the theory that meteors had their origin in lunar volcanoes.

Assuming that a lunar volcano may eject masses of matter with the requisite velocity to pass beyond the region where the lunar gravitation predominates over the terrestrial and the masses become obedient to the earth's attraction, Dr. Gould examines in detail the consequences to which the theory of the lunar origin of meteorites would necessarily lead, and presents his conclusions as follows:

"From the foregoing considerations we are warranted in assuming that for every body expelled from lunar volcanoes with a force adapted for throwing to the earth an aerolite of average dimensions there are, in probability, at the very least one hundred and eighty bodies expelled with forces not thus adapted; that for every mass ejected with the average force of a lunar volcano and striking the earth, there are at least one hundred and eighty masses of inadequate dimensions ejected; and that for any given combination of volcanic force and projected mass, the region of the lunar surface, whence the mass may reach the earth, is exceeded in extent by the tract of the lunar surface whence this would be impossible, in the ratio of fifty to one. Combining these several individual probabilities, it will readily be perceived that not more than

$$\frac{1}{50} \times \frac{1}{180} \times \frac{1}{180} = \frac{1}{1620000}$$

of all the ejected lava masses, or about 3 in 5,000,000 of each possible size, would probably ever reach the earth as aerolites; nor is this an unsafe estimate. It is a very guarded one, and the fraction $\frac{1}{2000000}$ would be more likely to be correct. Now, can we regard it as probable that the moon has parted with so large an amount of matter as nearly, if not quite, two million times the combined mass of all the aerolites which have fallen to the earth? I think not. Even of those known to have fallen there are more than five hundred of various weights, one of them having a mass of thirty thousand pounds. The tokens of such a mass of gravitative matter as this would imply could not fail to be

legibly inscribed in the unerring and enduring records of our system. They would preclude the accordance which is found to exist between the present lunar theory and the ancient observations. They would be found to be inconsistent with the known values of precession and nutation. They might, indeed, almost be said to be incompatible with the present mass of the moon."

From the observations of the meteor of November 15, 1859, *Prof. H. A. Newton*¹, of Yale College, concluded that it must have moved in a hyperbolic orbit, and that we have, therefore, two sources of meteors—the solar system and stellar space.

With regard to the periodic meteors of August, *Mr. A. C. Twining*², of New Haven, concluded that "the radiant is probably capable of a far more exact determination than is ordinarily supposed or than could have been anticipated, and it is apparently subject to a motion of several degrees from day to day, and a motion which exhibits some remarkable points of agreement in the comparison of one year's positions with those of other years."

From a discussion of the peculiar characteristics of the August meteors *Prof. H. A. Newton*³ came to the following conclusions:

- 1st. The individual meteors are cosmical bodies.
- 2d. They are permanent members of the solar system, revolving about the sun in elliptic orbits.
- 3d. The direction and velocity of the relative motion, and therefore of the absolute motion of the individual bodies, are nearly the same.
- 4th. The whole group forms what may be considered a ring or disk around the sun.
- 5th. The periodic time is two hundred and eighty-one days.

¹ A. J. S., XXX₂, 186.

² A. J. S., XXXII₂, 444.

³ A. J. S., XXXII₂, 448.

In the same paper Professor Newton estimates the whole number of meteors in the August ring as 300,000,000,000,000.

In March, 1862, *A. C. Twining*¹, published a paper entitled "Investigations respecting the phenomena of meteoric rings as affected by the earth," and arrived at the following conclusions: "The position of the node of the ring cannot be shifted by the earth's action more than a degree or two in half a million of years; there is an appreciable change of radiant positions, relative to locality on the earth's surface and to the hour of the day, whose maximum is about $3\frac{1}{4}^{\circ}$ between the extremes and to which the extremes approach; the terrestrial disturbance is sufficient to affect the perihelion distance of the meteors by many millions of miles and to expand the ring to a corresponding breadth at the ascending node; also to collect together in orbits, of similar elements, those meteors which are similarly affected in respect of radiant positions; and terrestrial disturbances do not appear sufficient to draw off meteors into permanently erratic orbits; so that, unless in exceptional instances, meteors are not lost to the ring other than those which the atmosphere absorbs or arrests. If meteors are partially arrested without being dissipated in an excessively tenuous upper medium it may be possible that the ordinary and unconformable meteors are such as have missed a return to the ring under the effect of atmospheric retardation."

Mr. Twining appends the suggestion that, "perhaps comets whose vastly extended atmospheres or heads around the nucleus, although greatly attenuated are perhaps competent to arrest meteors completely, may be found in rare instances to have been disturbed by *impact* with a meteoric ring whose mere attractive influence it would not be possible to detect."

In 1863, *Mr. B. V. Marsh*,² of Philadelphia, published a paper on "The luminosity of meteors as affected by latent heat," in which he arrived at the following results: "The upper regions of the atmosphere, even to its utmost limit,

¹ A. J. S., XXXIII., 244.

² A. J. S., XXXVI., 92.

are grand reservoirs of latent heat most admirably adapted to the protection of the earth from collision with bodies approaching it with planetary velocity from without. The intruder is instantly surrounded with a fiery envelope heated to the greatest conceivable intensity; its surface is burned off or dissipated into vapor; the sudden expansion of the stratum immediately beneath the burning surface tears the body into fragments, each of which, retaining its planetary velocity, is instantly surrounded by a similar envelope, which produces like effects, and so on until, in most cases, the whole is burned up or vaporized." A second paper on the same subject, and of similar import, was published by Mr. Marsh in the Proceedings of the American Philosophical Society, vol. XIV, 114.

From an examination of the list of November meteor showers from A. D. 902 to A. D. 1833 *Prof. H. A. Newton*¹ concluded that "the star-shower has a motion along the sidereal year of one day in seventy years, and also that the shower has a period of about a third of a century. This precession seems to imply that the orbit of the body furnishing these meteors has only a small inclination to the ecliptic, and that the motion is retrograde. The small distance of the radiant from the point to which the earth is moving, viz., 7°, confirms this conclusion."

In an article on the peculiarities of the November meteors, *Prof. H. A. Newton*² arrived at the following conclusions: "The length of the annual period as determined from the showers in A. D. 902 and 1833, reckoning 233 leap years, 19 odd days, and adding six hours for difference of longitude, is $365 + \left(\frac{233 + 19.25}{931}\right)$, or 365.271 days. The length of the cycle is 33.25 years.

"The length of the part of a cycle during which showers may be expected may be five or six years or, for extraordinary displays, at least 2.25 years. The supposition of a ring of uniform density throughout its circuit seems improbable.

¹ A. J. S., XXXVI., 300.

² A. J. S., XXXVIII., 53.

“The elements of the mean of the orbits of the different groups composing the partial ring are:

Semi-major axis = 0.98049,

Inclination = 17° ,

and the ring is nearly circular.

“The velocity with which these bodies enter the earth’s atmosphere is about 20.17 English miles per second.”

The most elaborate American paper on meteors up to the date of its publication was prepared in 1865 by *Prof. H. A. Newton*,¹ who discussed the subject under the following divisions, the conclusions being briefly stated in each case:

“1st. The average altitude of the middle points of the luminous portions of the meteor paths is found to be 59.4 English miles.

“2d. The relative frequency of meteors when the heavens were divided into eight equal parts was about equal in all—perhaps a slight preponderance in the southeast—and the relative frequency in different parts of the visible heavens may be considered a function of the zenith distance only.

“3d. Not quite *one* in fifty of all the meteors seen at any one place should have the middle points of their apparent paths within 10° of the zenith.

“4th. The number of visible meteors that come into the atmosphere every day would be 10,460 times the number visible at one station; or the average number that traverse the atmosphere daily, that are large enough to be seen with the naked eye, if the sun, moon, and clouds would permit, would be $30 \times 24 \times 10,460 = 7,531,000$.

“5th. The number of meteoroids in the space which the earth traverses is discussed at length and formulæ are derived for computing the whole number when the average number is known for a given unit of time. *

“6th. The average length of the apparent paths derived from 213 European and 803 American observations is $12^{\circ}.6$.

“7th. Adopting the theory that for every meteor visible to

¹ Mem. Nat. Acad. Sciences I, 291.

the naked eye there are 52.7 that are visible through a comet-seeker, the whole number of meteoroids coming daily into the air is 400,000,000.

"8th. The mean distance of the meteors from the observer is less than 144 miles.

"9th. The mean foreshortening of the meteor paths by perspective is from $16^{\circ}.0$ to $12^{\circ}.6$.

"10th. The average length of the visible part of meteor paths is between 24 and 40 or 21 and 34 miles; probably nearer 21 and 34 miles.

"11th. The mean duration of flight is not much, if any, greater than half a second of time."

*Prof. H. A. Newton*¹ discussed the observations of the altitudes of *seventy-eight* meteors observed on November 13-14, 1863, at Washington, Haverford College, Germantown, Philadelphia, and other points, giving diagrams exhibiting the altitudes of these meteors, and also of *thirty-nine* meteors observed in August, 1863, with the following results:

	November meteors.	August me- teors.
Mean altitude at appearance . .	96.2 miles.	69.9 miles.
Mean altitude at disappearance .	60.8 "	56.0 "
Mean altitude of middle point of path	78.5 "	62.9 "

In a paper on "The Theory of Meteors" *Prof. Daniel Kirkwood*² arrived at the following conclusions:

"The zodiacal light is probably a dense meteoric ring, or rather, perhaps, a number of rings.

"Variable and temporary stars are caused by the interposition of meteoric rings.

"Mercury's mean motion is probably diminished by the action of meteoric matter.

"The transit of a meteoric stream or cloud affords the most probable explanation of the phenomenon known as 'dark days.'

¹ A. J. S., XL₂, 250.

² Proc. A. A. A. S. 1866, 8.

"It seems probable that a ring of meteor asteroids exists within the orbit of Titan, Saturn's largest satellite, and causes the annual motion of the apsides of Titan, found by Bessel to be 30' 28".

"Saturn's rings are probably composed of an indefinite number of extremely minute asteroids or meteorites.

"The gaps in the distribution of the mean distances of the asteroids between Mars and Jupiter are analogous to the gaps in Saturn's rings."

In May, 1867, *Prof. Daniel Kirkwood* published a book¹ under the title "Meteoric Astronomy," designed by the author to present in a popular form the principal results of observation and study in that branch of Astronomy. It was devoted chiefly to the collection of some of the principal theories and the more important observations, and to presenting them in a brief but popular form without attempting to set forth any new theory.

A paper by *Prof. H. A. Newton*² in 1867, "On certain recent foreign contributions to Astro-meteorology," was devoted to the discussion of a table comparing the epochs and positions of radiant points of shooting-stars concluded independently by R. P. Greg and Dr. E. Heis; the influence of the August and November meteors on the temperature of the atmosphere; the paths and probable origin of the shooting-stars, by Schiaparelli, and the age of the November group of shooting-stars.

From the data obtained from the observations of the November meteors in 1867 *Prof. H. A. Newton*³ discussed the geographical limits of the shower; the personal equation of observers; the form of the curve of intensity; the breadth of the radiant in latitude; the length of the radiant in longitude, and the distribution in longitude of the perihelia of the orbits of the meteors.

¹ J. B. Lippincott & Co., Phila., 1867, 129 pp.

² A. J. S., XLIII₂, 285.

³ A. J. S., XLV₂, 89.

From data obtained from the observations of the November meteors of 1867 at the U. S. Naval Observatory and at Richmond, Va., *Prof. S. Newcomb*¹, U. S. N., computed the altitude of *nine* meteors, finding the mean altitude at apparition to be 102 miles, and at disappearance 47 miles. From data obtained from observations on the same occasion, *Prof. W. Harkness*², U. S. N., discussed a method of determining the mass of such meteors as are consumed before reaching the earth.

Assuming that the light produced is always proportional to the amount of material consumed, he arrived at the conclusion that "the mass of the ordinary shooting-stars does not differ greatly from one grain."

In 1869 a paper by *Prof. Daniel Kirkwood*³ on "Comets and Meteors" was devoted to exhibiting the probable coincidences between the orbits of comets and periodical meteors.

In 1871 *Mr. Jacob Ennis*⁴ published a paper entitled "The meteors and their long-enduring trails."

The scope and method of this paper are briefly sketched by the author, and are best presented in his own words, as follows:

"Firstly, I will bring forward many facts to prove that some meteors undergo a process of burning or oxidation while passing through the air, and that the trails are the smoke and ashes of such burning.

"Secondly, I will give facts and reasoning which show that some meteors are composed of various simple chemical elements unoxidized, and which are therefore capable of burning in the air.

"Thirdly, I will show the order and process of creation by which such meteors were originally formed and left in an unoxidized condition."

These points are discussed at length, and numerous theories and observations are cited as proof.

¹ A. J. S., XLV₂, 233.

² A. J. S., XLV₂, 237.

³ Proc. Am. Phil. Soc., XI, 215.

⁴ Proc. A. A. A. S. 1871, 122.

In a paper on the "Influence of meteoric showers on auroras" *Prof. Pliny E. Chase*¹ concludes that "there seems therefore good reason to look for an increase of auroral displays soon after every meteoric shower."

In discussing the meteors of November 27, 1872, *Prof. H. A. Newton*² remarked, "With Professor Weis and others, I am inclined to consider them all to have been once connected with periodic comets. The scattering took place apparently at or near the perihelion."

In 1872 *Prof. J. W. Mallet*, of the University of Virginia, read a paper³ on "The occluded gases of meteorites," and another paper⁴ by this author on the same subject appeared in 1875.

In 1875 *Prof. A. W. Wright*, of Yale College, published an account⁵ of some very carefully conducted experiments made to determine the character and quantity of the occluded gases of meteorites.

From these experiments he derived results differing materially from those obtained by other investigators.

This paper was followed by three others⁶ during 1875 and 1876, in which *Professor Wright* reached the conclusion that the spectra of gases from meteorites were identical with the spectra of comets.

In a lecture⁷ at the Sheffield Scientific School of Yale College, on "The relation of Meteorites and Comets," *Prof. H. A. Newton*, exhibiting a fragment from the meteoric stone which fell in Iowa February 12, 1875, very clearly presented his theory of the connection of these bodies.

The principal points in the theory, together with some of the arguments, may be briefly stated as follows:

Between the largest meteorite known and the faintest shooting-star that can be seen on a clear night with a telescope

¹ Proc. Am. Phil. Soc. XII, 401.

² A. J. S., V₃, 62.

³ Proc. Royal Society, XX, 365.

⁴ A. J. S., X₃, 206.

⁵ A. J. S., IX₃, 294.

⁶ A. J. S., X₃, 44; XI₃, 253; XII₃, 165.

⁷ Nature, Vol. XIX, 315, 340.

there is no essential difference as to astronomical character. In all their characteristic phenomena there is a regular gradation of meteors from one end of the line to the other. They differ in bigness, but in their astronomical relations we cannot divide them into groups. They are all similar members of the solar system. In proof of these statements we cite some of the points in which the large and small meteors are alike and unlike:

1st. They are all solid bodies. It is doubtful whether a small gaseous mass could exist permanently as a separate body in the solar system. A liquid would probably freeze and become solid. In any case, neither a gas nor a liquid could for an instant sustain the resisting pressure which a meteor is subjected to in the air, much less could it travel against it with the velocity observed in ordinary meteor flights. In short, every shooting-star must be a solid body.

2d. The large meteors and the small ones are seen at about the same height from the earth's surface. The air is a shield to protect the earth from an otherwise intolerable bombardment by these meteors. Some of the larger masses penetrate this shield, or, at least, are not melted before their final explosion, when the fragments, their velocity all gone, fall quietly to the ground. The small ones burn up altogether or are scattered into dust.

3d. The velocities of the large and the small meteors agree, and, though they are never measured directly very exactly, we are sure that in general they are more than two and less than forty miles per second. Velocities of from ten to forty miles per second imply that these masses are bodies that move about the sun as a center or else move through space. These velocities, as well as other facts, are utterly inconsistent with a permanent motion of such bodies about the earth or with a terrestrial or a lunar origin.

4th. The motions of the large and small meteors as they cross the sky have no special relations to the ecliptic. If either kind had special relations to the planets, in their origin or in their motions, we should have reason to expect them,

if not always, at least in general, to move across the sky *away from* the ecliptic. The fact is otherwise. Both large and small meteors are seen moving *towards* the ecliptic as often as *from* it. Neither class seem, therefore, to have any relation to the planets.

Again, in general character the two classes are alike. They have like varieties of color; they have similar luminous trains behind them. In short, we cannot draw any line dividing the stone or iron producing meteor from the shooting-star, at least in their astronomical relations. They are all astronomically alike. They differ in size; but that has nothing to do with their motion about the sun or in space.

The general connection between comets and meteors may be exhibited in the peculiar relations existing between the meteors of November 13-14 and their accompanying comet. The orbit of these meteors is one that is described in 33.25 years. The meteors go out a little further than the planet Uranus, or about twenty times as far as the earth is from the sun. While they all describe nearly the same orbit they are not collected in one compact group. On the contrary, they take four or five years to pass a given place in the orbit, and are to be thought of as a train several hundred millions of miles long but only a few thousands of miles in thickness.

Along with this train of meteors travels a comet. It passed the place where we meet the meteor stream nearly a year before the great shower of 1866 and two or three years before the quite considerable displays of 1867 and 1868. How came it that this comet and the meteors travel the same road? The plane of the comet's orbit might have cut the earth's orbit to correspond with any other day of the year than November 15; or, cutting it at this place, the comet might have gone nearer to the sun or farther away; or, satisfying these two conditions, it might have made any angle from 0° to 180° instead of 467° ; or, satisfying all these, it might have had any other periodic time than 33.25 years; even then it might have gone off in any other direction of the plane than that in which the meteoroids were

traveling. All these things did not happen by chance; there is something common.

The comet alluded to is not the only one that has an orbit common with meteors, though it is the only case in which the orbit of the meteors is *completely* known, aside from our knowledge of that of the comet. Every August, about the tenth day, we have an unusual number of meteors—a star-sprinkle as it has been called. A comet whose period is about 125 years moves in the plane and probably in a like orbit with these meteors. Near the first of December there have been several star-showers, notably one in 1872, and these meteors are traveling nearly in the orbit of Biela's comet. In April, too, some showers have occurred which are thought to have had something to do with a known comet. Thus much as to the meteors of the star-showers. The sporadic meteors are with good reason presumed to be (and observed facts *prove* some of them to be) the outliers of a large number of meteor streams.

Considering again the November meteor stream and its comet we find that the several bodies move along a common path not at all by reason of a present physical connection. They are too far apart—in general, a thousand times too far apart—to act on each other so much that we may measure the effect. Their connection has been in the past. They must have had some common history. Looking now at the comets, we see that they have been apparently growing smaller at successive returns. Halley's comet was much brighter in its earlier than in its later approaches to the sun. Biela's comet has divided into two or more principal parts, and seems to have entirely gone to pieces. Several comets have had double or multiple nuclei. In the year 1366, in the week after the star-shower, a comet crossed the sky exactly in the track of the meteors. A second comet followed in the same path a week after. Both belonged, no doubt, to the November stream, and one of them may perhaps have been the comet of 1866.

The November meteor stream is a long, thin one. We

have crossed the stream at many places along a length of a thousand millions of miles, sometimes in advance of and sometimes behind the comet, and all along this length have been found fragments—sometimes few, sometimes many. This form of the stream suggests continuous action producing it. A brief, violent action *might* have given this form, but a slowly acting cause seems more natural.

Again, in the history of Biela's comet we have distinct evidence of continued action. The comet divided into two parts not long before 1845, and yet in 1798 fragments of it were met with so far from the comet that they must have left the comet long before, probably many centuries ago.

"Thus we are led to say, *first*, that the periodic meteors of November, of August, of April, &c., are caused by solid fragments of certain known or unknown comets coming into our air; *secondly*, that the sporadic meteors, such as we can see any clear night, are the like fragments of other comets; *thirdly*, that the large fire-balls are only larger fragments of the same kind; and, *finally*, that a portion broken off from one of those large fragments in coming through the air must once have been *a part of a comet*."

"How came the comet to break up? Perhaps the prior question would be, How came the comet together? In its history there is much that cannot yet be explained, much about which we can only speculate."

"Thus, how came this meteoric stone to have its curious interior structure? As a mineral it resembles more the deepest fire-rocks than it does the outer crust of the earth. It seems to have been formed in some large mass, possibly in one larger than any of our existing comets. Some facts show that the comets have almost surely come to us from the stellar spaces. Out somewhere in the cold of space a condensing mass furnished heat for the making of this stone. The surrounding atmosphere was unlike ours, since some of these minerals could hardly have been made in the presence of the oxygen of our air. Either in cooling or by some catastrophe the rocky mass may have been broken to pieces,

so as to enter the solar system having little or no cohesion, like a mass of pebbles; or it may have come, and probably did come, a single solid stone.

In either case, as it got near to the sun new and strong forces acted on it.

The same heat and repulsion that develops and drives off from a comet in one direction a tail, sometimes a hundred millions of miles long, may have cracked off and scattered in another direction solid fragments. One of these contained in it this stone, and it wandered in its own orbit about the sun, itself an infinitesimal comet, how many thousands of millions of years we know not, until three years ago it came crashing through the air to the earth in Iowa."

More than ordinary space has been given to the citations from the various statements and arguments and to the concluding speculation of Professor Newton's paper because, better than any preceding American discussion, it presents the status of the modern theories of meteors and comets which are now generally accepted by the scientific world.

The latest formal discussion of this subject was presented by *Professor Newton*¹ in his presidential address before the American Association for the Advancement of Science, at Buffalo, in 1886. This address was devoted wholly to the consideration of the various theories in regard to the motions, character, and origin of meteorites, meteors, and shooting-stars.

The discussion in this address follows the same general lines as in the lecture just cited, while the various arguments are presented with far greater elaboration. No new hypotheses or theories are offered; but the key-note of the address, given in the author's own words, is that "science may be advanced by rejecting bad hypotheses as well as by forming good ones."

¹ Proc. Am. Ass. Ad. Science 1886, 1.

EXAMINATION OF THEORIES.

The abstracts and excerpts just presented are, from the limitations of the method employed, frequently very brief, sometimes disconnected, and generally separated from the various discussions which led to the results cited.

Although they present in themselves insufficient data for an accurate study or a rigorous discussion of the subject, they are quite sufficient to illustrate the evolution of the modern theories as they have been successively developed from the superstitions and the dogmatic assumptions of the last century.

This development is a fair illustration of the growth of most of the sciences, and the sometimes absurd and baseless theories, some of which have been cited, are the usual evidences of an anxious, persistent searching after the truth which is satisfied only by success.

While the modern theories have been slowly evolved from a multitude of observations and discussions, expanding here and there along the lines of least difficulty, it is not improbable that frequently there has been a lack of the nicest discrimination as to what were real and well-established facts.

Keeping in view the precept that no sound theory can be based on doubtful data, it is proposed to examine briefly the accumulated mass of so-called knowledge of Meteors and Comets, with a view to ascertaining what we actually *know* about these bodies; what we *infer*, *assume*, and *assert*, and to some extent, perhaps, what we *do not know* about them.

METEORS.

Those bodies which are usually designated as *meteors*, *meteorites*, and *shooting-stars* are known, to some extent, by every intelligent person. The first name is usually applied to those sporadic bodies which one can see occasionally on any clear night; the second term is applied to iron or stony masses that sometimes fall to the earth, while the last term is used

to designate those bodies which appear in such periodic showers as those of November 13-14, August 6-10, etc., but which, like the first named, are, almost without exception, entirely consumed before they reach the earth.

These bodies have received, at various times, a great variety of names, such as "Fiery Tears of St. Lawrence," "Fire-balls," "Bolides," "Aerolites," "Meteoroids," etc., most of which have been coined to suit the fancy or ambition of some aspiring author. The only definite knowledge we have of this class of bodies before they reach the surface of the earth is obtained with the spectroscope, and the results from observations with that instrument indicate that all these bodies are similar in composition, and their spectra are the same as that obtained from those masses that have reached the surface of the earth before destruction.

There appears to be, therefore, no reason for using but two names—the one, *meteor*, for those bodies that are consumed before they reach the earth; and the other, *meteorite*, for the solid iron or stony substances that succeed in storming our atmospheric barriers, reaching the surface of the earth intact and bringing our only material messages from the depths beyond.

Sporadic meteors as well as meteorites move apparently in all directions. Meteors that appear in showers seem to emanate from pretty well defined points in the heavens, each separate shower having its own radiant, and in most cases the bodies are not condensed in a single compact mass, but are scattered along the orbit in which they move.

This orbit has been determined for several of the showers with considerable accuracy.

From the testimony of the meteors themselves nothing is known of their origin. The theories of a terrestrial or a lunar volcanic origin are easily shown to be absurd, while the so-called theories that place their origin in other portions of the solar system are mere idle speculations.

COMETS.

The whole number of comets, real and suspected, from about 1770 B. C. to the end of 1889 A. D., the elements of whose orbits have *not* been computed, is 472. From 370 B. C. to the end of 1889 A. D. the number of comets the elements of whose orbits have been computed is 309. Of these, 18 are known to have *elliptic* orbits. In the case of 52, the computed elliptic orbits have not been verified by observation.

The computations show that 231 have *parabolic* orbits, and indicate that 7 have *hyperbolic* orbits.

Thus it appears that not more than seven per cent. of the comets whose orbits have been discussed are known to have *elliptic* orbits, while it is almost certain that seventy-five per cent. have *parabolic* orbits. Of course, the periodic comets, whatever their origin, belong now to the solar system.

As it is highly improbable that there are two or more kinds of comets of intrinsically diverse character and of different origin, it follows that all the comets had their genesis beyond the limits of the solar system, and that the few periodic comets are the exception to the general law, and at best are only adopted members of the solar family.

There are only two sources of actual knowledge of the physical constitution of comets :

One is from the use of the spectroscope; the other is the behavior of the light from a star when seen through various portions, but especially the nucleus of a comet.

As is well known, observations with the spectroscope are not always easily interpreted, but in this case the difficulty is not so great as at first it seems to be.

It is a general law that where there is a continuous spectrum containing all the primary colors without gaps, the light is derived from an incandescent solid or liquid body. A discontinuous spectrum containing bands or bright lines indicates that the light comes from luminous gases or vapors.

To these general rules there are some important exceptions or modifications.

If the temperature of certain vapors or gases be raised to a high degree the number and the appearance of the colored band or of the bright lines change rapidly, though not uniformly, and some investigators have asserted that if the temperature be raised to something over 4,500° Fah. the spectrum will become practically continuous. Similar changes in the phenomena are observed if a gas, like hydrogen, is rendered luminous by the electric spark and then subjected to varying pressures. With a pressure amounting to one-twentieth of an inch of mercury the spectrum is discontinuous and consists of several groups of bright lines in the green. As the pressure is gradually increased there appears a temporary spectrum of bands, then a spectrum of three lines, afterwards a more permanent and complete spectrum of bands, and finally, under a pressure of 52 inches of mercury, a complete and pure continuous spectrum.

The spectra of comets, which have been obtained by careful and experienced observers, present a large number of variations and combinations, ranging from one or more faint bands with indistinct or fluted borders against a colorless background to a faint continuous spectrum with bands or lines of a greater or less degree of brightness and definition.

The most obvious interpretation of the spectroscopic observations of comets is that the bands and lines are the true spectra of a gaseous body, varying through a wide range under the effect of changing pressure and temperature, superimposed upon the faint continuous spectrum derived from the sunlight reflected from the nucleus or other parts of the comet.

Such is the information derived from the spectroscope.

In the vast number of observations of comets, made for the determination of their positions or their physical peculiarities, it has sometimes been noted that the comet passed

between the observer and a star without diminishing the apparent brightness of the star or changing its position.

While observing Comet I, 1866, in January, 1866, I saw on one occasion the nucleus of the comet pass directly over a star of the 9.2 magnitude with no more effect on the brightness of the star than would be produced by the close proximity of any object as bright as the comet's nucleus. Similar accounts have been given by other observers, and the phenomenon is too well attested to admit of a reasonable doubt.

The light from a star could not pass unobstructed through a solid body or a dense aggregation of solid bodies; and, considering this phenomenon alone or in connection with the appearance of the nucleus as it approaches and recedes from perihelion, it appears that we are driven to the conclusion that the nucleus of a comet is composed principally, if not entirely, of gaseous matter, which varies in form and in density from the effect of the sun's attraction and repulsion.

COMETS AND METEORS.

The elements of the orbits of *four* meteor streams have been determined with considerable accuracy. These are the streams that produce the showers of November 13-14, November 27, April 20, and August 10.

It has also been found that the orbit of the meteor stream of November 13-14 coincides very closely with the orbit of Comet I, 1866. The orbit of the November 27 stream corresponds to that of Biela's comet, the orbit of the April stream to that of comet I, 1861, and the orbit of the August stream is nearly identical with that of Comet III, 1862.

The identify of these orbits is quite as good as could be expected from the uncertain character of the observations on which the adopted positions of the meteor streams depend.

On these coincidences in the orbits of meteor streams and of certain comets depends principally the modern theory of comets and meteors, which, briefly stated, is as follows:

Sporadic meteors, individual members of meteoric showers,

and meteorites differ in magnitude and appear under widely varying conditions, but from an astronomical standpoint they are all alike.

They are all solid bodies and are fragments of comets.

Assuming that this theory is true, we shall find that some of the inferences drawn from it are of great importance in their bearing on cosmical physics.

1st. As the meteoric masses, both great and small, are derived from comets, they must have originated beyond the limits of the solar system, and they furnish evidence of the existence in space of exactly such minerals, though in different combinations, as are found in the earth's crust.

2d. They arise from the disintegration of comets, which for centuries have furnished the material for the enormous areas of bodies forming the various meteor streams that trail along the orbits of these masses for immense distances.

3d. The meteors forming the shower of November 13-14 have been observed for more than 900 years, and yet the comet whose gradual destruction has produced these bodies was not discovered until 1866. The August meteors have been observed for more than six centuries, but the comet whose disintegration has furnished the material for this vast stream remains intact, and was not discovered until 1862.

The accepted *comet-meteor* theory does not explain clearly the visibility of comets or the changes that occur in the apparent brightness and in the density of the nucleus as these bodies approach and recede from the sun; neither does it explain in a satisfactory manner the position of the comets in their attendant meteor streams. If comets are composed of solid matter or of discrete solid particles, it would seem quite proper to ask why they become visible at such immense distances from the earth and the sun.

The perihelion distance of 26 per cent. of the comets with known orbits is equal to or greater than the mean distance of the earth from the sun.

Many comets when first seen are much farther from the sun than is the earth at aphelion, and the spectroscope only

gives the information that the light is derived from a gas or vapor. From our constant experience with solid masses of stone and iron on the surface of the earth and under the unobstructed influence of the sun, it is impossible to see how the sun's heat alone can produce gas or vapor from such bodies at the observed distances.

As the comet approaches the sun the faint diffused mass of the body begins to contract, and a point in the mass, generally nearer the sun than the center, becomes brighter and denser, frequently, as it rapidly nears the sun, changing its form and brightness in a marked manner from day to day.

It is not improbable that the solid constituents of meteorites would be vaporized if they passed as near the sun as did Comet II, 1882; but it is not probable that this change does occur at distances greater than the radius of the earth's orbit, if it is effected simply by the action of the sun.

If the visibility is caused by the assumed enormous change of temperature experienced by the solid portion of the comet in passing from outer space to the locus of visibility in the solar system, then the entire mass of the comet should be vaporized and solid meteoric bodies would cease to exist.

If, on the other hand, this visibility is brought about by the effect of this change of temperature on the occluded gases stored up in the solid portions of the comet, then during the long period in which these masses are subjected to the solar action these gases would all be expelled and dissipated and none would be found in those meteorites which finally find their way to the surface of the earth and into the chemist's laboratory.

The meteors of the shower of November 27 are scattered along the orbit of that stream for at least 500 millions of miles. If this elongation of the meteor stream is formed, as is highly probable, by the difference in velocity between those meteors on that portion nearest the sun and those on the outside of the mass, then, if the comet is the meteor-producing body, the same action would tend to break it up and destroy it early in its existence as a solar satellite.

If the existence of the comet as a member of the solar system antedates the meteor stream, it is difficult to see how the comet could have remained intact long enough to have been observed, in the presence of forces that for thousands of years have been transforming the figure of the original mass and stretching it out into a stream whose length is measured by hundreds of millions of miles. It is not improbable that comets of large dimensions are destroyed by the action of such forces; but that a body of that character should miraculously survive its own destruction and be found existing in ordinary cometary form in the midst of its own ruins is a proposition that makes large demands on the imagination.

If the brightness of comets is caused by the vaporization of iron or stony matter, it must be produced by collisions between the masses at such velocities that a high temperature is developed, producing an incandescent vapor yielding a distinctive spectrum. It seems difficult to explain how such relative velocities can arise among the individual members of the same stream moving in a common orbit. It is more than probable that the light of a star passing from the extremely low temperature of space through the supposed high temperature of the comet's nucleus, and again into the temperature of space, would suffer so much apparent change of position that it would compel recognition. It is claimed, however, that the individual masses of meteoric matter which form the nucleus are so far separated that the light of a star can pass through the aggregated mass without material change of direction.

But if the masses are vaporized by collisions, then there must be absolute contact, which would to a great extent obstruct the passage of stellar light and would be certain to produce refraction.

Lockyer's Theories.—before leaving the consideration of these points I venture to call attention for a moment to a recent theory which has been set forth with considerable elaboration of detail.

In a paper entitled "Researches on the Spectra of Meteorites," presented to the Royal Society on October 4, 1887, and in the Bakerian lecture on April 12, 1888, followed by an appendix to the same on January 10, 1889, Mr. Lockyer presented in detail his laboratory experiments, combined with the more or less accurate observations of other astronomers and physicists, which led him to certain definite conclusions in regard to the relations of comets and meteors.

The author's conclusions and theories can be most succinctly presented in the following citations from the papers mentioned:

"The existing distinction between stars, comets, and nebulae rests on no physical basis."

"All self-luminous bodies in the celestial spaces are composed of meteorites or masses of meteoric vapor produced by heat brought about by condensation of meteor swarms due to gravity."

"Meteorites are formed by the condensation of vapors thrown off by collisions. The small particles increase by fusion brought about again by collisions, and this increase may go on until the meteorites may be large enough to be smashed by collisions when the heat of impact is not sufficient to produce volatilization of the whole mass."

"Beginning with meteorites of average composition, the extreme forms, iron and stony, would in time be produced as the result of collisions."

"The spectra of all such bodies depend upon the heat of meteorites produced by collisions and the average space between the meteorites in the swarm, or, in the case of consolidated swarms, upon the time which has elapsed since complete vaporization."

"The temperature of vapors produced by collisions in nebulae, stars without C and F, but with other bright lines, and in comets away from perihelion is about that of the Bunsen burner."

"The temperature of the vapors produced by collisions in α Orionis and similar stars is about that of the Bessemer flame."

"The brilliancy of the aggregated masses depends upon the number of the meteorites and not upon the intensity of the light."

"The bright flutings of carbon in the spectra of some 'stars,' taken in conjunction with their absorption phenomena, indicate that widely separated meteorites at a low temperature are involved."

"New stars are produced by the clash of meteor-swarms, the bright lines seen being low temperature lines of those elements in meteorites the spectra of which are most brilliant at a low stage of heat."

"A comet is a swarm of meteors in company. Such a swarm finally makes a continuous orbit by virtue of arrested velocities. Impacts will break up large stones and will produce new vapors, which will condense into small meteoroids."

"When the meteorites are *strongly* heated in a glow-tube the whole tube, when the electric current is passing, gives us the spectrum of carbon. When a meteor-swarm approaches the sun the whole region of space occupied by the meteorites * * * gives us the same spectrum."

"The first stage in the spectrum of a comet is that in which there is only the radiation of the magnesium. The next is that in which Mg. 500 is replaced wholly or partially by the spectrum of cool carbon. Mg. is then added and cool carbon is replaced by hot carbon. The radiation of manganese 558 and sometimes lead 546 is then added. Absorption phenomena next appears, manganese 558 and lead 546 being indicated by thin masking effect upon the citron band of carbon. The absorption band of iron is also sometimes present at this stage. At this stage also the group of carbon flutings, which I have called carbon B, probably also makes its appearance. As the temperature increases still further, magnesium is represented by *b*, and lines of iron appear. This takes place when the comet is at or near perihelion."

"The observations on meteorites recorded in the Bakerian

Lecture and the discussion of cometary observations contained in this Appendix show that the vapors which are given out by the meteorites as the sun is approached are in an approximate order: slight hydrogen, slight carbon compounds, magnesium, sodium, manganese, lead, and iron. Now, of these the hydrogen and carbon compounds are alone permanent gases, and the idea is that they have been occluded as such by the meteorites."

"The aurora being a low temperature phenomenon, we should expect to find in its spectrum lines and remnants of flutings seen in the spectra of meteorites at low temperatures. The characteristic line of the aurora is the remnant of the brightest manganese fluting at 558."

"The spectrum of the nebulae, except in some cases, is associated with a certain amount of continuous spectrum, and meteorites glowing at a low temperature would be competent to give the continuous spectrum with its highest intensity in the yellow part of the spectrum."

"Only seven lines in all have been recorded up to the present in the spectra of nebulae, three of which coincide with lines in the spectrum of hydrogen and three correspond to lines in magnesium. The magnesium lines represented are the ultra-violet low-temperature line at 373, the line at 470, and the remnant of the magnesium fluting at 500, the brightest part of the spectrum at the temperature of the Bunsen burner. The hydrogen lines are *h*, *F*, and *H γ* . (434). Sometimes the 500 line is seen alone, but it is generally associated with *F* and a line at 495. The remaining lines do not all appear in one nebulae, but are associated one by one with the other three lines."

"When a tube is used in experiments to determine the spectrum of meteoric dust at the lowest temperature we find that the dust in many cases gives a spectrum containing the magnesium fluting at 500, which is characteristic of the nebulae and is often seen alone in them. If the difference between nebulae and comets is merely of cosmographical position, one being out of the solar system and one being in

it; and, further, if the conditions as regards rest are the same, the spectrum should be the same, and we ought to find this line in the spectrum of comets when the swarm most approaches the undisturbed nebulous condition, the number of collisions being at or near a minimum—*i. e.*, when the comet is near aphelion the fluting should be visible alone.”

After citing the results of the spectroscopic observations of several comets, the author remarks: “This spectroscopic evidence is of the strongest, but it does not stand alone. Comets at aphelion present the telescopic appearance, for the most part, of globular nebulae.”

The comprehensive theory set forth in the quotations just cited assumes that the auroræ, nebulae, meteorites, comets, and most of the stars all have a common origin, and that all the multifarious telescopic and spectroscopic phenomena exhibited by these bodies are due to the varying velocities of the collisions between the meteoric particles and masses of which in some form all these bodies are composed.

We are told that meteorites at a low temperature present in the spectrum a certain line, at 558, due to manganese, and also that this line appears in the nebulae, the aurora, and in comets at considerable distances from perihelion. Hence the identity of all these bodies is inferred and the foundation of the theory is laid.

Meteorites are subjected to laboratory experiments in tubes in which the temperature is gradually raised to a high degree and the varying spectra is noted. Spectroscopic observations of nebulae, comets, and stars are then compiled and classified, until the several groups are so arranged that they present nearly the same sequence of spectra that have been derived from meteoric matter at increasing temperatures in the experiments.

The theory is then extended and we are given to understand that when, in the case of nebulae and stars greater activity of collisions occur, or when a comet approaches the sun, the same phenomena appear and in the same order.

The identity of these bodies is then supposed to be complete and the theory established.

This theory of collisions rests upon a remarkable congeries of experiments, observations, and assumptions. Many of the observations and many of the laboratory experiments, which were made by the author, as well as much of the data quoted throughout his papers are entitled to the highest merit. But, considering much of the data and many of the statements in his conclusions, and especially the extraordinary assertion that "comets at aphelion present the telescopic appearance for the most part of globular nebulae," it is not remarkable to find the author's data, as well as his deductions, vigorously attached by able physicists.

Huggins.— After a careful study of the spectrum of the aurora *Mr. Huggins*¹ remarks: "After consideration, I think that I ought to point out that *Mr. Lockyer*'s recent statement that 'the characteristic line of the aurora is the remnant of the brightest manganese fluting at 558' is clearly inadmissible, considering the evidence we have of the position of this line."

After a very thorough study of the spectra of the nebulae, *Mr. Huggins*² writes: "As, therefore, there seems to be little doubt that the 'remnant of the fluting at 500' is not coincident with the brightest nebular line, and the next most characteristic group of this spectrum, the triplet at 3720, 3724, and 3730, according to *Living* and *Dewar*, does not appear to be present in the photographs, we may conclude that the remarkable spectrum of the gaseous nebulae has not been produced by burning magnesium."

*Professor Living*³ says in regard to the line denoted by *Lockyer* as 470: "I have never seen the line at λ 4703 in the spectrum of the magnesium flame. As it is a conspicuous line in the arc and spark, we looked for it in the flame, but did not find it."

¹ Proc. Roy. Soc., XLV, 435.

² Proc. Roy. Soc., XLVI, 55.

³ Proc. Roy. Soc., XLVI, 56.

If the testimony of Huggins and of Liveing and Dewar represents the observed phenomena, and their observations have not yet been disproved, then most of the broad theories of Lockyer, which assume a common origin and structure for auroræ, nebulæ, comets, and stars, lacks a basis of observed facts, resting wholly, so far as the aurora and nebulæ are concerned, on approximate coincidences in the spectra, while the assumed telescopic appearance of comets at aphelion is a creation of the imagination.

CONCLUSIONS.

Attention has been called to these various theories relating to comets and meteors simply with a view to emphasizing the fact that none of the systems, whether simple or complex, seems to explain all the observed phenomena.

As a scientific explanation, the direct and simple is always preferable to the indirect and involved method, and this safe precept should be the guide in all investigations of the apparent physical connection between comets and meteors.

It seems to me that the true theory of the origin and the relations of comets and meteors is yet to be discovered.

When asked to give my own theory of these bodies I can only reply that I have none. At the same time I see less objection to the following hypotheses than to any of those now doing duty as theories:

Meteors and meteorites are solid iron or stony bodies and, whatever their origin, are now members of the solar system. Comets are composed chiefly of gaseous matter, and originate outside of the solar system. Some of these bodies on entering the sphere of solar attraction are so far drawn away from their original orbits by the masses of the sun's outer satellites that they become permanent members of the solar system. Of these, at least *four* have become entangled in the immense aggregations known as meteor streams and have adopted the orbits of their captors. The meteors still remain meteors, however, and the comets retain their former identity and peculiar structure.

OBSERVATIONS OF METEORS.

Most of the observers of sporadic meteors and meteorites have been either amateurs or persons entirely deficient in that special training which is so essential in a trustworthy observer of unexpected phenomena. Fortunately, however, most of the important phenomena have been noted by intelligent and skilled observers, whose zeal and care have left little to be desired.

It would be impracticable to mention even the names of all the successful observers, but any sketch of the progress of meteoric astronomy in this country would be notably deficient if some of the prominent names were omitted.

The remarkable meteor shower of November 13, 1833, attracted the attention of many careful observers and zealous students along our Atlantic coast, and for several years the subject was carefully investigated by Prof. Dennison Olmstead and Prof. A. C. Twining, who were the pioneers in the study of this science in the United States.

From 1838 when E. C. Herrick began his work he labored with untiring industry as an observer and a compiler of observations and other data until his death, in 1862, and no one in this country did so much as he in promoting the observation and investigation of the August meteors.

Mr. Herrick also gave considerable attention to the study and observation of the November meteors, but this stream was made a special study by Prof. H. A. Newton, with the best results.

Professor Newton's observations of the November meteors began in 1860 and have been continued to the present time, while his investigations of the motions and character of this stream place him undeniably at the head of American workers in this branch of Astronomy.

Much work of the highest value was done by Prof. C. U. Shepard and by Prof. J. Lawrence Smith in the chemical examination of all classes of meteorites, and excellent in-

vestigations of a similar character have been carried out by other eminent chemists in the country.

The zeal and industry of Professor Shepard was shown in his extensive collection of meteoric specimens, which at the time of his death was the largest in America.

The attempt to bring together all the published observations in this country in one systematic collection is a task beset with grave difficulties.

The reports of these observations are scattered through all the scientific journals, the metropolitan and local newspapers, and the proceedings of all grades of learned societies. Frequently the reports, when found, have but little scientific value from lack of the necessary information. In many instances much time and space are wasted in describing trivial details which have no interest or value in connection with the true meteoric phenomena, while the really essential data are not mentioned.

It sometimes happens that the only available information in regard to a meteorite is derived from the report of its chemical examination, and there can be found no astronomical data whatever to account for its position; it is simply a portion of the earth's surface, and the *how*, *when*, and *whence* of its advent remain unanswered.

It has been impossible, sometimes, to find any trustworthy authority for essential data, and it has been necessary frequently to interpret freely where the observer or writer has given but a slight clue to his meaning.

In nearly all cases marginal references are made to the original papers in order to facilitate further examination, if desired.

THE CATALOGUES.

The catalogues of Sporadic Meteors, Meteoric Showers, Observed Meteorites, and Discovered Meteorites are supposed to contain all observations, accompanied with the necessary data that have been found in the various publications to which the author has had access. It is not assumed, how-

ever, that these lists contain all the good observations that have been made in this country ; in fact, it is quite certain that they do not, and one of the principal aims of this paper will be attained if this fact attracts sufficient attention to bring to light the missing or the unpublished observations.

In all the catalogues the *day* of the observation is the *astronomical* day. It was manifestly impracticable to give every reference to each object in the five catalogues, and only the most important ones have been retained.

Occasionally references are only given to the first page of a paper when it contains several observations of the same phenomenon.

In the reference notes at the bottom of the page the principal abbreviated notation may be explained as follows :

A. J. S., XXV., 306, refers to the American Journal of Science, Vol. XXV, *second* series, page 306.

Trans. A. P. S. refers to the American Philosophical Society.

Proc. A. P. S. refers to the American Philosophical Society.

Proc. A. A. A. S. refers to the proceedings of the American Association for the Advancement of Science.

CATALOGUES

I.-V.

CATALOGUE I.

Number.	DATE.				Locality.	Iron or stone.
	Year.	Month.	Day.	Hour. Min.		
1	1781(?)				Portage Bay, Chilcot Inlet, Alaska	I.
2	1807	December	13	18.5	Weston, Conn.	S.
3	1810	January	30		Caswell, N. C.	S.
4	1823	August	7		Nobleborough, Me.	S.
5	1825	February	10		Nanjemoy, Md.	S.
6	1827	May	9		Sumner Co., Tenn.	S.
7	1828	June	4		Richmond, Va.	S.
8	1829	May	8	3	Forsyth Co., Ga.	S.
9	1829	August	15		Deal, N. J.	S.
10	1835	July	30	2	Charlotte, Dickson Co., Tenn.	I.
11	1837	May	5	3	East Bridgewater, Mass.	S.
12	1839	February	13	3	Little Piney, Pulaski Co., Mo.	S.
13	1840	October			Concord, N. H.	S.
14	1843	March	25		Bishopville, S. C.	S.
15	1844	January			Argentine Republic	I.
16	1846	August	14	3	Cape Girardeau, Mo.	S.
17	1847	February	25	3	Marion, Linn Co., Iowa	S.
18	1848	May	19	16	Castine, Me.	S.
19	1849	October	31	3	Charlotte, Cabarrus Co., N. C.	S.
20	1855	August	5	3	Lincoln Co., Tenn.	S.
21	1857	April	1		Costa Rica, Central America	S.
22	1857				Independence Co., Iowa	S.
23	1859	March	28	4	Harrison Co., Ind.	S.
24	1859	July	4		Crawford Co., Ark.	S.
25	1859	August	11		Bethlehem, N. Y.	S.
26	1860	May	1	1	New Concord, Ohio	S.
27	1865	March	24	21	Vernon Co., Wis.	S.
28	1868	November	27	5	Danville, Ala.	S.
29	1868	December	5	3	Frankfort, Ala.	S.
30	1869	October	5	23	Stewart Co., Ga.	S.
31	1871	May	20	20	Searsmont, Me.	S.
32	1874	May	14	2.5	Nash Co., N. C.	S.
33	1875	February	12	10.5	Iowa Co., Iowa	S.
34	1876	June	24	21	Kansas City, Mo.	I.
35	1876	December	21	8.75	Rochester, Fulton Co., Ind.	S.
36	1877	January	2		Warrenton, Warren Co., Mo.	S.
37	1877	January	23		Cynthiana, Harrison Co., Ky.	S.
38	1879	May	10	5	Estherville, Emmet Co., Iowa	S. & A.
39	1879	August			Fomatlan, Jalisco, Mexico	
40	1883				Calderilla, Chill	I.
41	1885	November	27		Mazapil, Mexico	I.
42	1886	March	27	3	Johnson Co., Ark.	I.
43	1887	January	21	2	De Cewsville, Haldimand Co., Ontario	S.
44	1890	May	2	5 15	Winnebago Co., Iowa	S.

1. Cat. State Mining Bureau of Cal., 1888, No. 2925.

2. { Trans. Amer. Phil. Soc. VI₁, 323.
A. J. S. XXXVII₁, 130; VI₂, 410.3. A. J. S. II₂, 392.4. A. J. S. VII₁, 170; IX₁, 400.5. A. J. S. IX₁, 351; X₁, 131; VI₂, 406.6. A. J. S. XVII₁, 326; XVIII₁, 200, 378.7. A. J. S. XV₁, 195; XVI₁, 191.8. A. J. S. XVIII₁, 388.

9. Proc. A. A. A. S., 1851, Vol. II, 188.

10. A. J. S. XLIX₁, 336.11. A. J. S. XXXII₁, 395.12. A. J. S. XXXVII₁, 385; XXXIX₁, 254.13. A. J. S. IV₂, 353; VI₂, 416.14. A. J. S. II₂, 392; VI₂, 411.

15. Proc. Lit. and Phil. Soc. Liverpool, VII, 1853.

16. A. J. S. XXXII₃, 229.17. A. J. S. IV₂, 288, 429.18. A. J. S. VI₂, 251, 406.19. A. J. S. IX₂, 143; X₂, 127.20. A. J. S. XXIV₂, 134; XXXI₂, 264.

21. Buchner, 93.

22. A. J. S. XXX₂, 208.

Observed Meteorites.

Number.	Weight.	Authority.	Remarks.
1	88 lbs.	Seen to fall by the father of one of the oldest Indians.
2	300 "	Nathan Wheeler.....	Observed by many persons.
3	3 "	Madison.	
4	4 "	A. Dinsmoor.	
5	16.5 "	W. D. Harrison.	
6	11 "		
7	4 "		
8	36 "	Elias Beall.	
9			
10	9 lbs.	G. Troost.....	Weight "rather more than half an ounce."
11	0.5 lb.		J. L. Smith gives the date as August 1.
12	50 lbs.	Mr. Harrison.	
13			Weight 370.5 grains.
14	13 lbs.	C. U. Shepard.	
15		H. E. Symonds.....	Fall witnessed by 1,400 soldiers. About a cubic yard of the mass remained above the surface of the ground.
16	4.5 lbs.	E. S. Dana.	
17	130 lbs.	S. L. Penfield.	
18	0.1 lb.	D. C. Rogers.	
19	19.5 lbs.	Giles Gardner.	
20	3.9 "	H. Bost.	
21		James B. Dooley.	
22		C. U. Shepard.....	Fell in the "summer" of 1857.
23	3.7 lbs.		Several observers.
24		Mr. Scott.	
25		C. U. Shepard	"Smaller than a pigeon's egg."
26	460.2 lbs.		
27		J. L. Smith.	
28	4.5 lbs.	W. Brown.	
29	1.7 "	Jas. W. Hooper.	
30	0.8 lb.	Mrs. Buck.	
31	12 lbs.		
32			
33	500 lbs.	Many observers.
34			
35	0.8 lb.	A. J. Morris.	
36	100 lbs.		
37	15 "	"Fell in the afternoon."
38	750 "		
39		C. U. Shepard	Several pieces; the largest weighed about 2 lbs.
40	"Small."	Ward and Howell.....	Not yet described.
41	8.7 lbs.	W. E. Hidden.	
42	107.5 "	G. F. Kunz.	
43	0.75 lb.	E. E. Howell.	
44	184 lbs.	G. F. Kunz.....	Probably more fragments to be discovered.

23. A. J. S. XXVIII₂, 409.
24. Owens' 2d Geolog. Reconnaissance of Arkansas, 408.
25. A. J. S. XXX₂, 206.
26. { A. J. S. XXX₂, 103, 207, 296;
A. J. S. XXX₁, 87; XXXII₂, 30.
27. A. J. S. XII₃, 207.
28. A. J. S. XLIX₂, 90.
29. A. J. S. XLVII₂, 240.
30. A. J. S. L₂, 335, 339.
31. A. J. S. II₃, 133, 200.
32. A. J. S. X₃, 147.
33. A. J. S. IX₂, 407, 459; X₃, 44, 357.
34. A. J. S. XI₃, 316.
35. A. J. S. XIII₃, 207, 243.
36. A. J. S. XIII₃, 243; XIV₃, 219.
37. A. J. S. XIII₃, 243; XIV₃, 219.
38. A. J. S. XVII₃, 77, 186; XIX₃, 459, 495; XX₃, 136.
39. A. J. S. XXX₃, 105.
- 40.
41. A. J. S. XXXIII₃, 221.
42. A. J. S. XXXIII₃, 494, 500.
43. Science, N. Y., March 7, 1890, 167.
44. Science, N. Y., May 16, 1890, 304.

Number.	DATE.		Locality.	Iron or stone.
	Year.	Month and day.		
1	1735		Arizona.....	I.
2	1784		Bahia, Brazil.....	I.
3	1792		Zacatecas, Mexico.....	I.
4	1808		Red River, Texas.....	I.
5	1810		Santa Rosa, New Granada.....	I.
6	1811		Durango, Mexico.....	I.
7	1818		Lockport, N. Y.....	I.
8	1819		Burlington, N. Y.....	I.
9	1820		Guilford Co., N. C.....	I.
10	1822		Randolph Co., N. C.....	I.
11	1826-7		Waterloo, Seneca Co., N. Y.....	I.
12	1828		Bedford Co., Pa.....	I.
13	1832		Walker Co., Ala.....	I.
14	1834		Scriba, Oswego Co., N. Y.....	I.
15	1834		Claiborne, Clark Co., Ala.....	I.
16	1835		Buncombe Co., N. C.....	I.
17	1836		Brazos, Texas.....	I.
18	1839	March.....	Putnam Co., Ga.....	I.
19	1839		Buncombe Co., N. C.....	I.
20	1840	February 26.....	Chili.....	I.
21	1840		Cooke Co., Tenn.....	I.
22	1840		Smithland, Livingston Co., Ky.....	I.
23	1841	February.....	Lexington Co., S. C.....	I.
24	1842		Grayson Co., Va.....	I.
25	1842		Roanoke Co., Va.....	I.
26	1842		Carthage, Tenn.....	I.
27	1842		Green Co., Tenn.....	I.
28	1845		DeKalb Co., Tenn.....	I.
29	1845		Otsego Co., N. Y.....	I.
30	1846		Franconia, N. H.....	I.
31	1846		Jackson Co., Tenn.....	I.
32	1847		Chesterville, S. C.....	I.
33	1847-8		Murfreesborough, Tenn.....	I.
34	1849		Pittsburgh, Pa.....	I.
35	1850		Allegheny Co., Pa.....	I.
36	1850		Seneca River, N. Y.....	I.
37	1850		Salt River, Ky.....	I.
38	18 0		Botetourt Co., Va.....	I.
39	1853		Jefferson Co., Tenn.....	I.
40	1853		Union Co., Ga.....	I.
41	1853	July.....	Campbell Co., Tenn.....	I.
42	1853	August.....	Tazewell, Claiborne Co., Tenn.....	I.
43	1854		Madoc, Ontario.....	I.
44	1854		Haywood Co., N. C.....	I.
45	1855		Coahuila, Mexico.....	I.
46	1856		Nelson Co., Ky.....	I.
47	1856		Nebraska.....	I.
48	1856		Madison Co., N. C.....	I.
49	1856		Forsyth, Taney Co., Mo.....	I.
50	1856		Marshall Co., Ky.....	I.
51	1856		Denton Co., Texas.....	I.
52	1856		Oktibbeha, Miss.....	I.
53	1857		Laurens Co., S. C.....	I.
54	1858		Washington Co., Wis.....	I.

1. { Smithsonian Report 1863, 55, 85.
A. J. S. XVIII₂, 369; XIX₂, 161, 162.
2. { A. J. S. XV₂, 12; XXXVI₃, 158.
Sci. Am. Supp. Oct. 19, 1889.
3. A. J. S. XV₂, 11.
4. A. J. S. VII₁, 218; XVI₁, 217; XXVII₁, 382.
5. A. J. S. XV₂, 11.
6. A. J. S. XV₂, 19.
7. A. J. S. XLVIII₁, 388; II₂, 374, 391.
8. A. J. S. XLVI₁, 401; II₂, 391; XV₂, 20.
9. A. J. S. XVII₁, 140; XL₁, 369.
10. A. J. S. XVII₁, 140; II₂, 391; XV₂, 21.
11. A. J. S. XI₂, 39; XXXIV₂, 298.
12. A. J. S. II₂, 391; XV₂, 21.
13. A. J. S. XLIX₁, 344; II₂, 391; XV₂, 21.
14. A. J. S. XL₁, 366; II₂, 390; IV₂, 75.
15. A. J. S. XXXIV₁, 332; XLVIII₁, 145.
16. A. J. S. XXXVI₁, 81; XLIII₁, 359.
17. A. J. S. XXXI₂, 459.
18. A. J. S. XVII₂, 331.
19. A. J. S. IV₂, 82.
20. Lon., Ed. and Dublin Phil. Mag. X₄, 12.
21. A. J. S. XXXVIII₁, 250; XLIII₁, 351.
22. A. J. S. II₂, 357; XV₂, 21.
23. { A. J. S. X₂, 128; XV₂, 5, 16.
Proc. A. A. A. S. 1850, Vol. I, 152; 1853, Vol. II, 189.
24. A. J. S. XLIII₁, 169; II₂, 392.
25. A. J. S. XLIII₁, 169; II₂, 392.
26. A. J. S. II₂, 356; XV₂, 20.
27. A. J. S. XLIX₁, 342; II₂, 391.

Discovered Meteorites..

Number.	Weight.	Authority.	Remarks.
1	1,400 lbs.	Joseph Henry.....	The "Ainsa" or "Tucson" meteorite.
2	11,819 "	The "Bendego" meteorite.
3	2,000 "		
4	1,635 "		
5	1,700 "		
6		Weight 30,000 or 40,000 lbs.
7	36 lbs.	B. Silliman.	
8	160 "	B. Silliman.	
9	28 "	C. U. Shepard.	
10	2 "	C. U. Shepard.	
11	0.1 lb.	C. U. Shepard.....	Rammelsburg thought this was not a meteorite.
12	C. U. Shepard.....	"A few ounces." Dr. Genth thought it was Spiegeleisen.
13	165 lbs.	G. Troost.	
14	8 "	C. U. Shepard.	
15	40 "	C. T. Jackson.	
16	30 "	C. U. Shepard.....	One date given is 1845.
17	324 "	C. U. Shepard.	
18	72 "	J. E. Willett.	
19	1.3 "	C. U. Shepard.	
20	17 "	R. P. Greg.....	Found in the desert of Tarapaca.
21	2,000 "	G. Troost.	
22	9.9 "	G. Troost.	
23	117 "	C. U. Shepard.....	Found on "Ruff's Mountain."
24	J. B. Rogers.....	"Original mass of many pounds' weight."
25	W. B. Rogers.....	No weight given.
26	280 lbs.	G. Troost.	
27	29 "	G. Troost.....	Found near Babb's mill.
28	36 "	G. Troost.	
29	C. U. Shepard.....	Weighed 276 grains.
30	Robert Gilmore....	Weighed "15 or 20 pounds."
31	G. Troost.....	Weighed 15 oz.
32	36 lbs.	C. U. Shepard.	
33	19 "	G. Troost.	
34	0.6 lb.	F. A. Genth.	
35	292 lbs.	B. Silliman.	
36	9 "	Date somewhat doubtful.
37	8 "	B. Silliman.	
38	C. U. Shepard.....	Original mass lost.
39	2.5 lbs.		
40	15 "		
41	0.3 lb.		
42	60 lbs.	C. U. Shepard.....	Also described by J. L. Smith.
43	370 "	T. Sterry Hunt.	
44	C. U. Shepard.....	Weight about $\frac{1}{8}$ oz.
45	252 lbs.	J. L. Smith.....	The "Couch" meteorite.
46	0.21 lb.	G. J. Brush.	
47	35 lbs.		
48	0.11 lb.	G. J. Brush.	
49	197 lbs.	C. U. Shepard.	
50	15 "	C. U. Shepard.	
51	C. U. Shepard.....	Weighed 66 grains.
52	0.3 lbs.	W. J. Taylor.....	Found in an Indian mound.
53	4.7 "	W. E. Hidden.	
54	85.8 "	F. Breundecke.	

28. A. J. S. XLIX₁, 341; II₂, 391.
29. A. J. S. II₂, 391; XV₂, 16.
30. A. J. S. II₂, 392; IV₂, 87.
31. A. J. S. II₂, 357; XV₂, 21.
32. A. J. S. VII₂, 449; XV₂, 21.
33. A. J. S. V₂, 351; XV₂, 21.
34. A. J. S. XV₂, 22; XI₃, 72.
35. A. J. S. XV₂, 7; Proc. A. A. A. S. 1850, Vol. II, 37.
36. A. J. S. XIV₂, 439; XV₂, 363.
37. A. J. S. XV₂, 22; Proc. A. A. A. S. 1850, 36.
38. A. J. S. XLI₂, 250.
39. A. J. S. XVII₂, 329.
40. A. J. S. XVII₂, 328.
41. A. J. S. XIX₂, 153.

42. A. J. S. XVII₂, 131, 325; XIX₂, 153.
43. A. J. S. XIX₂, 417.
44. A. J. S. XVI₂, 327.
45. { A. J. S. XIX₂, 160.
Smithsonian Report, 1863, 56.
46. A. J. S. XXX₂, 240; XXXI₂, 459.
47. A. J. S. XXX₂, 204; XXXII₂, 146.
48. A. J. S. XXX₂, 240; XXXI₂, 459.
49. A. J. S. XXX₂, 205; XXXIV₂, 467.
50. A. J. S. XXX₂, 240; XXXI₂, 459.
51. { A. J. S. XXXI₂, 459.
Trans. St. Louis Acad. Sci. I, 623.
52. A. J. S. XXIV₂, 293.
53. A. J. S. XXXI₂, 463.
54. Smithsonian Report 1869, 417.

CATALOGUE II.—Dis-

Number.	DATE.		Locality.	Iron or stone.
	Year.	Month and day.		
55	1858-9	Augusta Co., Va.....	I.
56	1859	Rogue River Mountains, Ore.....	I.
57	1860	Mountains of East Tennessee.....	I.
58	1860	Franklin Co., Ky.....	I.
59	1860	October	La Grange, Oldham Co., Ky.....	I.
60	1860	December.....	Coopertown, Tenn.....	I.
61	1860	Newton Co., Ark.....	I.
62	1863	Ren-selaer Co., N. Y.....	S.
63	1863	February 18	Colorado.....	I.
64	1863	Tucson, Arizona.....	I.
65	1863	June 9.....	Dakota.....	I.
66	1866	Bear Creek, Colo.....	I.
67	1866	Frankfort, Ky.....	I.
68	1867	Allen County, Ky.....	I.
69	1868	April	Losttown, Cherokee Co., Ga.....	I.
70	1868	"Southeastern Missouri".....	I.
71	1868	Auburn, Macon Co., Ala.....	I.
72	1869	Utah.....	S.
73	1869	El Dorado Co., Cal.....	I.
74	1869	Trenton, Wis.....	I.
75	1870	Howard Co., Ind.....	I.
76	1873	August.....	Madison Co., N. C.....	I.
77	1873	Cleburne Co., Ala.....	I.
78	1874	Wacanda, Kan.....	S.
79	1875	San Francisco, Brazil.....	I.
80	1877	Whitfield Co., Ga.....	I.
81	1878	Fayette Co., Texas.....	S.
82	1879	Whitfield Co., Ga.....	I.
83	1879	July 19.....	Davidson Co., N. C.....	I.
84	1879	Ivanpah, Cal.....	I.
85	1880	Eagle Station, Carroll Co., Ky.....	I.
86	1880	Lexington Co., S. C.....	I.
87	1880	Rutherford Co., N. C.....	I.
88	1882	June 10.....	Maverick Co., Texas.....	I.
89	1882	Burke Co., N. C.....	I.
90	1883	May 15.....	Grand Rapids, Mich.....	I.
91	1883	Little Miami Valley, Ohio.....	I.
92	1883	Wayne Co., W. Va.....	I.
93	1884	June.....	Independence Co., Ark.....	I.
94	1884	Hammond, St. Croix Co., Wis.....	I.
95	1884	August.....	Santa Fé County, New Mexico.....	I.
96	1884	Chili.....	I.
97	1885	Catorze, San Luis Potosi, Mex.....	I.
98	1887	January	Laramie Co., Wyoming.....	I.
99	1887	March	Claiborne Co., Tenn.....	I.
100	1887	March	Cumberland Co., Tenn.....	S.
101	1887	March 27.....	Chattooga Co., Ga.....	I.
102	1888	April 30.....	Welland, Ontario.....	I.
103	1888	Chili.....	I.
104	1888	Chili.....	I. & S.
105	1888	Chili.....	I.
106	1888	Hamilton Co., Texas.....	I.

55. A. J. S. XV₃, 337.

56. Proc. Boston Soc. Nat. Hist. Vol. 7, 161, 174, 175, 191, 279, 289.

57. { A. J. S. XXXIV₃, 473.

{ Proc. Acad. Nat. Sci. Phil. 1886, 366.

58. Smith. Report 1868, 343; A. J. S. XLIX₃, 331.59. A. J. S. XXXI₂, 151, 265.60. A. J. S. XXXI₂, 151, 266.61. A. J. S. XL₂, 213; XXXIV₃, 471.62. A. J. S. XXXIV₃, 60.63. A. J. S. XLII₂, 218.64. { A. J. S. XXXVI₂, 152.

{ Proc. Cal. Acad. Sci. III, 30.

65. A. J. S. XXXVI₂, 259.66. A. J. S. XLII₂, 250, 286; XLIII₂, 280.67. A. J. S. XLIX₃, 331.68. A. J. S. XXXII₃, 500.69. A. J. S. XLVI₂, 257; XLVII₂, 234.70. A. J. S. XLVII₂, 233.71. A. J. S. XLVII₂, 230.72. A. J. S. XXXII₃, 226.73. A. J. S. III₃, 438; VI₃, 18.74. A. J. S. XLVII₂, 271; III₃, 69.75. A. J. S. V₃, 155; VII₃, 391.76. A. J. S. XII₃, 439.77. A. J. S. XIX₃, 370; XX₃, 74.78. A. J. S. XI₃, 473; XIII₃, 211.

79. { Comptes Rendus LXXXIII, 917, 918; LXXXIV, 478, 482, 1085, 1508.

{ A. J. S. XXIII₃, 232; XXIX₃, 33, 496.

covered Meteorites—Cont'd.

Number.	Weight.	Authority.	Remarks.
55	152 lbs.	J. W. Mallet.	
56	John Evans	Mass above ground, 4.5 x 3.5 feet.
57	254 lbs.	F. A. Genth.	
58	0.1 lb.	G. J. Brush.	
59	112 lbs.	Found by Wm. Daring.
60	37 "	Found by D. Crockett.
61	J. L. Smith.....	Specimen weighed 22½ oz.
62	3.3 lbs.	S. C. H. Bailey.	
63	29 "	Found near Central City by Otho Curtice.
64	632 "	
65	10.6 "	Dr. Jackson	Found in the "Dakota Indian country."
66	436 "	J. L. Wilson.	
67	24 "	J. L. Smith.	
68	24.3 "	J. E. Whitfield.	
69	6.6 "	
70	0.8 lb.	C. U. Shepard.	
71	8 lbs.	C. U. Shepard.	
72	1.9 "	E. S. Dana.	
73	85 "	B. Silliman.	
74	143.5 "	J. L. Smith.....	Six fragments found; the first in 1869.
75	4 "	E. T. Cox.	
76	25 "	B. S. Burton.	
77	35.8 "	W. E. Hidden.	
78	116 "	C. U. Shepard.	
79	22,048 "	E. Guignet.....	In the province of San Catherina.
80	13 "	W. E. Hidden.	
81	321 "	Whitfield and Merrill.	
82	117 "	C. U. Shepard.	
83	2.8 "	W. E. Hidden.	
84	128.2 "	C. U. Shepard.	
85	80 "	G. F. Kunz.	
86	10.5 "	C. U. Shepard.	
87	4.8 "	L. G. Eakins.	
88	97.25 "	W. E. Hidden.	
89	1 lb.	G. F. Kunz.	
90	114 lbs.	J. R. Eastman.	
91	G. F. Kunz.....	Fragments found in mounds by F. W. Putnam; now in the Peabody Museum.
92	26 lbs.	G. F. Kunz.....	Several fragments.
93	94 "	W. E. Hidden.	
94	53 "	Davenport Fisher.	
95	324.4 "	G. F. Kunz.....	Several fragments.
96	14.5 "	Ward and Howell.	Found near Puquios. Not yet described.
97	92 "	G. F. Kunz.	
98	25.06 "	G. F. Kunz.	
99	18 "	G. F. Kunz.	
100	94.5 "	J. E. Whitfield.	
101	27 "	G. F. Kunz.	
102	17.5 "	E. E. Howell.	
103	16 "	Ward and Howell.	Thirty leagues east of Taltal. Not yet described.
104	27 "	Ward and Howell.	Thirty-five leagues S. E. of Taltal. Not yet described.
105	Ward and Howell.	Estimated at from 6 to 8 lbs.
106	179 "	Ward and Howell.	Found five miles south of Carlton. Not yet described.

80. A. J. S. XIV₃, 246; XXI₃, 286.
 81. A. J. S. XXXVI₃, 113.
 82. A. J. S. XXVI₃, 336; XXXIV₃, 473.
 83. A. J. S. XX₃, 324.
 84. A. J. S. XIX₃, 381.
 85. A. J. S. XXXIII₃, 228.
 86. A. J. S. XXI₃, 117.
 87. A. J. S. XXXIX₃, 395.
 88. A. J. S. XXXII₃, 304; XXXIII₃, 115.
 89. A. J. S. XXXVI₃, 275.
 90. A. J. S. XXVII₃, 299; XXX₃, 312.
 91. A. J. S. XXXIII₃, 228.
 92. A. J. S. XXXI₃, 145; Proc. A. A. S., 1885, 246.

93. A. J. S. XXXI₃, 460; School of Mines Quarterly, Columbia Coll., VII, No. 2, 188.
 94. A. J. S. XXXIV₃, 381.
 95. A. J. S. XXX₃, 235; XXXII₃, 311.
 96.
 97. A. J. S. XXXIII₃, 233.
 98. A. J. S. XXXVI₃, 276.
 99. A. J. S. XXXIV₃, 475.
 100. A. J. S. XXXIV₃, 387, 476.
 101. A. J. S. XXXIV₃, 471.
 102. Science, N. Y., March 7, 1890, 167.
 103.
 104.
 105.
 106.

CATALOGUE III.—Discovered

Number.	Locality.	Iron or stone.	Weight.
1	Tucuman, Argentine Republic.....	I.	1 lb.
2	Crawford Co., Arkansas.....	I.	30,000 lbs.
3	British America.....	S.	1.4 "
4	Canyon City, Trinity Co., Cal.	I.	386 "
5	Los Angeles, Cal.....	I.	19 "
6	San Bernardino Co., Cal.....	I.	80 "
7	Chihuahua.....	S.	
8	Atacama, Chili.....	I.	3,853 lbs.
9	Sierra de Chaco, Chili.....	I.	
10	Durango, Mexico.....	S. & I.	
11	Oaxaca, Mexico.....	I.	72.75 "
12	San Luis Potosi, Mexico.....	I.	
13	Xiquipilco, Mexico.....	I.	108.6 "
14	Mexico.....	I.	192 "
15	Mexico.....	I.	
16	Mexico.....	I.	
17	Mexico.....	I.	
18	Mexico.....	I.	2,942 lbs.
19	Mexico.....	I.	0.5 lb.
20	Mexico.....	I.	
21	Mexico.....	I.	5,000 lbs.
22	Bates Co., Missouri.....	I.	90 "
23	Ironton, Missouri.....	I.	
24	Rasgata, New Granada.....	I.	129 lbs.
25	Rockingham Co., N. C.....	I.	11 "
26	Green Co., Tenn.....	I.	6 "
27	Jefferson Co., Tenn.....	I.	290 "
28	Augusta Co., Va.....	I.	1.4 "
29	Augusta Co., Va.....	S.	5.5 "
30	Augusta Co., Va.....	I.	3.5 "
31	Augusta Co., Va.....	I.	36 "
32	Augusta Co., Va.....	I.	56 "
33	Augusta Co., Va.....	I.	2.2 "
34	Kiowa Co., Kansas.....	I.	
35	Chili.....	I.	95.5 lbs.
36	Jackson Co., Oregon.....	I.	2 oz.

1. A. J. S. XXXIV₃, 59.
2. A. J. S. XV₂, 12.
3. Owen's 2d Geological Reconnaissance of Ark., 408.
4. Trans. Roy. Soc. Canada, IV, sect. III, 97.
5. A. J. S., XXIX₃, 469.
6. A. J. S. IV₃, 495.
7. A. J. S. XXXV₃, 490.
8. { A. J. S. XIX₂, 163; II₃, 335; III₃, 207.
Proc. A. A. A. S. 1871, 266.
9. Buchner, 127.
10. Buchner, 131.
11. A. J. S. XXXVII₃, 439.
12. A. J. S. XV₂, 21.
13. Buchner, 149.
14. A. J. S. XV₂, 20; XXII₂, 374; XXIV₂, 295.
15. A. J. S. XXIX₃, 232.
16. Proc. A. A. A. S. 1871, 269.
17. Proc. A. A. A. S. 1871, 269.

Meteorites without Date.

Number.	Authority.	Remarks.
1	R. B. Riggs.....	Found in Col. Abert's collection of minerals presented to the National Museum.
2		
3		
4	A. P. Coleman	Brought to Coburg, Canada, in 1869.
5	C. U. Shepard.....	
6	C. T. Jackson.....	
7	G. P. Merrill.....	Put through an ore-crusher before its character was known.
8	J. L. Smith.....	
9		
10	J. E. Whitfield.....	
11		
12		
13		
14	Numerous pieces.
15	N. T. Lupton.....	
16	J. L. Smith.....	The San Gregorio meteorite, 6.5 feet long, 5.5 wide, 4.0 high.
17	J. L. Smith.....	"The largest yet found in that vicinity."
18	J. L. Smith.....	The "Butcher" meteorites—six, weighing 290, 430, 438, 550, 580, and 654 lbs.
19	J. L. Smith.....	Found in a collection of minerals from Mexico.
20	Now in National Museum, Washington, D. C. Estimated weight, 2,500 lbs.
21	W. M. Pierson.....	
22	G. C. Brodhead.....	
23	Small specimen.
24		
25	J. L. Smith.....	
26	C. U. Shepard.....	Property of the Smithsonian Institution; place of discovery unknown.
27	W. P. Blake.....	
28	R. B. Riggs.....	Riggs considers this a doubtful specimen.
29	L. G. Eakins.....	Probably found in Texas.
30	J. W. Mallet.....	
31	J. W. Mallet.....	
32	J. W. Mallet.....	
33	G. F. Kunz and J. W. Mallet.....	
34	F. H. Snow	Found by "cowboys" before that portion of Kansas was settled; fragments weighing in the aggregate more than 1,600 lbs. discovered.
35	Ward and Howell.....	Found about 10 or 12 leagues east of the port of Chafialar; not yet described.
36	Ward and Howell.....	Piece of a mass found by a miner.

18. Proc. A. A. A. S. 1871, 269.

19. A. J. S. XLV₂, 77.

20.

21. Smithsonian Report, 1873, 419.

22. A. J. S. X₃, 401; XII₁₃, 213.

23.

24. A. J. S. XV₂, 11.

25. A. J. S. XII₁₃, 213.

26. A. J. S. XXI₁₃, 119.

27. A. J. S. XXXI₃, 41.

28. A. J. S. XXXIV₃, 60.

29. A. J. S. XXXIX₃, 59.

30. A. J. S. II₃, 10.

31. A. J. S. II₃, 10.

32. A. J. S. II₃, 10.

33. A. J. S. XXXIII₃, 58.

34. Science N. Y., Vol. XV, No. 379, 290; No. 384, 359

35.

36.

CATALOGUE IV.—

Ref. number.	DATE.			TIME OF SHOWER.		Time of max. flight.	Whole number counted.	Max. hourly rate.	Radiant point.	No. of observ'g.
	Year.	Month.	Day.	Beginning.	End.					
				<i>h. m.</i>	<i>h. m.</i>	<i>h. m.</i>				
1	1803	April...	19	13 0	15 0			800		
2	1833	Nov....	13	9 0	Sunrise	16 0	200,000		"Bend of the Sickle" in Leo	
3	1834	"	12	14 50	17 25	17 0	18			1
4	1834	"	12				Many.		"In Leo"	
5	1835	"	13							
6	1835	"	13							
7	1835	"	13							
8	1836	"	12	14 0	16 0		248			
9	1836	"	12	15 0	18 15		253			
10	1836	"	12	15 30	16 0	15 40	29			
11	1836	"	12				23			1
12	1836	"	12				75			
13	1836	"	12				500			
14	1836	"	12				18			
15	1837	Aug.	9	8 0	15 0				$\alpha = 55^\circ, \delta = +60^\circ$	1
16	1837	Nov....	12	13 5	Sunrise	16 30	226		Near μ Leonis	8
17	1837	"	12	14 0	"		70		In Leo	1
18	1837	"	12	14 0	18 0		45		"	1
19	1837	"	12	14 45	16 37		74			12
20	1837	"	12	15 15	17 15	16 0	34	17		1
21	1837	"	12	15 45	17 0		52		In Leo	1
22	1838	April...	20	10 0	16 0		154		"	2
23	1838	Aug.	8	11 30	12 30		20		$\alpha = 55^\circ, \delta = +60^\circ$	1
24	1838	"	9	9 30	16 20	16 15	54		$\alpha = 35^\circ, \delta = +69^\circ$	1
25	1838	"	9	15 0	15 45		24			1
26	1838	"	10	8 55	10 0		36		In Cassiopeæ	3
27	1838	"	10	9 0	11 0		48			2
28	1838	"	10	9 30	16 0		122		Near ϵ Cassiopeæ	1
29	1838	"	10	14 30	16 0		65		"Sword handle of Perseus"	2
30	1838	Nov.	11	12 0	18 0	15 30	199	53	In Leo	8
31	1838	"	12	12 0	17 0	13 30	131	37	"	8
32	1838	"	13	13 0	17 20	15 30	233	80	"	8
33	1838	"	22	13 20	15 30	14 30	50	25		8
34	1838	Dec....	6	8 0	17 0		113		In Perseus	5
35	1838	"	7	8 0	11 0		210		"	5
36	1838	"	7	10 0	11 0		78		"Zenith"	3
37	1838	"	8	7 15	9 0		59			3
38	1838	"	11	8 45	10 0		18			1
39	1838	"	12	6 0	13 30		28			3
40	1838	"	15	6 0	9 15		9			4
41	1839	April...	18	12 0	15 0		58		$\alpha = 273^\circ, \delta = +45^\circ$	2
42	1839	"	18	14 0	16 0		25			2
43	1839	"	19	12 15	13 15		19			4
44	1839	"	19	14 0	16 0		25			2
45	1839	Aug.	4	9 0	11 0		36		Vicinity of Cassiopeæ	2
46	1839	"	9	9 7	14 7	13 30	691	194	Sword handle of Perseus	4
47	1839	"	9	12 41	15 36		100			1
48	1839	"	10	9 0	16 0		187	66		1
49	1839	"	10	9 0			76			2
50	1839	"	10	9 30	10 0		33		"Algenib"	1

1. A. J. S. XXXVI, 358; Va. Gazette and General Advertiser, April 23, 1803; N. H. Gazette, May 31, 1803; Med. Repository, N. Y., Vol. 1, 1804.
2. A. J. S. XXV, 354, 363.
3. A. J. S. XXVII, 335; XXVIII, 305.
4. A. J. S. XXVII, 339.
5. A. J. S. XXX, 375.
6. " " " "
7. A. J. S. XXX, 376.
8. A. J. S. XXXI, 390.
9. A. J. S. XXXI, 388.
10. A. J. S. XXXI, 390.
11. A. J. S. XXXI, 389.

12. A. J. S. XXXI, 391.
13. A. J. S. XXXI, 392.
14. A. J. S. XXXI, 392.
15. A. J. S. XXXIII, 133.
16. A. J. S. XXXIII, 379.
17. " " " "
18. " " " "
19. " " " "
20. " " " "
21. " " " "
22. A. J. S. XXXIV, 398.
23. A. J. S. XXXV, 167.
24. " " " "

Meteor Showers.

Ref. number.	Authority.	Place of observation.	Remarks.
1	Richmond, Va., and Ports-	"Fell too fast to be counted."
2	Prof. D. Olmstead.....	mouth, N. H. New Haven, Conn.....	Number estimated; seen throughout the Atlantic and Gulf States.
3	Prof. A. D. Bache.....	Philadelphia, Pa.	Observed on the morning of the 13th.
4	Prof. A. C. Twining.....	West Point, N. Y.....	"Unusual number;" hourly rate estimated at 40 or 50.
5	Amenia, N. Y.	"Unusual number." Observed on the morning of the 14th.
6	Mt. St. Mary's College, Md..	"Unusual number."
7	Salisbury, N. C.....	Many seen that were not recorded.
8	New York, N. Y.....	Cloudy.
9	S. Dunster.....	Springvale, Me.	Observed one hour.
10	New Haven, Conn.....	Whole number estimated at 400.
11	Cambridge, Mass.....	
12	Newark, N. J.....	
13	Randolph and Macon Col-	
		lege, Va.	
14	J. L. Russell.....	Hingham, Mass.	
15	G. C. Schaeffer.....	New York, N. Y.....	Counted between 200 and 300.
16	Prof. D. Olmstead	New Haven, Conn.....	Majority left trains.
17	G. C. Schaeffer.....	New York, N. Y.....	Many trains.
18	Prof. F. A. P. Barnard.	New York, N. Y.	
19	Prof. E. Loomis.....	Hudson, Ohio.	
20	E. Fitch.....	New Haven, Conn.	
21	L. Obermeyer	Mt. St. Mary's College, Md.	
22	Prof. Wright.....	Knoxville, Tenn.	
23	G. C. Schaeffer.....	Barren Hill, Pa.	
24	T. R. Dutton.....	Wilmington Island, Ga.	
25	W. A. Sparks.....	Society Hill, S. C.	
26	J. D. Dana.....	Norfolk, Va.	
27	E. C. Herrick.....	New Haven, Conn.	
28	T. R. Dutton.....	Wilmington Island, Ga.	
29	C. G. Forshey.....	Rock Island, Ill.	
30	Prof. J. Lovering.....	Cambridge, Mass.	Only two observers after 17 hours.
31	Prof. J. Lovering.....	Cambridge, Mass.	
32	Prof. J. Lovering.....	Cambridge, Mass.	
33	Prof. J. Lovering.....	Cambridge, Mass.	
34	E. C. Herrick.....	New Haven, Conn.	
35	E. C. Herrick.....	New Haven, Conn.	
36	Prof. A. W. Smith.....	Middletown, Conn.	
37	E. C. Herrick.....	New Haven, Conn.	
38	E. C. Herrick.....	New Haven, Conn.	
39	E. C. Herrick.....	New Haven, Conn.	
40	E. C. Herrick.....	New Haven, Conn.	
41	E. C. Herrick.....	New Haven, Conn.	
42	Prof. E. Loomis.....	Hudson, Ohio.	
43	E. C. Herrick.....	New Haven, Conn.	
44	Prof. E. Loomis.....	Hudson, Ohio.	
45	E. C. Herrick.....	New Haven, Conn.	
46	E. C. Herrick.....	New Haven, Conn.	
47	T. R. Dutton.....	Columbia, Tenn.	
48	L. L. Knox.....	Middletown, Conn.	
49	Charles Baldwin.....	New York, N. Y.	
50	Prof. A. W. Smith	Middletown, Conn.	

25. A. J. S. XXXV₁, 167.
 26. " " " "
 27. " " " "
 28. " " " "
 29. Trans. Am. Phil. Soc. VII, 266.
 30. A. J. S. XXXV₁, 323.
 31. " " " "
 32. " " " "
 33. " " " "
 34. A. J. S. XXXV₁, 361; XXXVI₁, 355.
 35. A. J. S. XXXV₁, 361.
 36. " " " "
 37. " " " "

38. A. J. S. XXXV₁, 361.
 39. " " " "
 40. " " " "
 41. A. J. S. XXXVI₁, 361.
 42. " " " "
 43. " " " "
 44. " " " "
 45. A. J. S. XXXVII₁, 325.
 46. A. J. S. XXXVII₁, 325.
 47. A. J. S. XXXVIII₁, 203.
 48. A. J. S. XXXVI₁, 325.
 49. " " " "
 50. " " " "

CATALOGUE IV.—

Ref. number.	DATE.			TIME OF SHOWER.		Time of max. flight.	Whole number counted.	Max. hourly rate.	Radiant point.	No. of observ's.
	Year.	Month.	Day.	Begin-ning.	End.					
				<i>h. m.</i>	<i>h. m.</i>	<i>h. m.</i>				
51	1839	Aug. ...	10	10 0	13 0	13 0	491	189	Sword handle of Perseus.....	4
52	1839	" ...	10	11 20	12 20		50		" "	1
53	1839	" ...	11	9 0	10 0		28		" "	3
54	1839	" ...	14	10 0	12 0		72	44	Ret. Cassiopeæ and Perseus..	3
55	1840	" ...	9	8 0	16 0		171		Sword handle of Perseus.....	1
56	1840	" ...	9	10 0	15 30	15 0	818	332	" "	4
57	1840	" ...	9	14 15	15 45		112	76	$\alpha = 30^\circ, \delta = +63^\circ 30'$	1
58	1840	" ...	10	15 0	15 35		35		" "	1
59	1841	April... 18	8 30	11 30			60		$\alpha = 198^\circ, \delta = -8^\circ$	1
60	1841	" ... 19	11 0	13 0			20		" "	3
61	1841	Aug. ... 9	12 0	15 0			49		" "	1
62	1841	" ... 10	9 0	10 0			60		" "	2
63	1842	April... 20	10 20	16 0	15 30		151	55	Corona Borealis	5
64	1842	Aug. ... 8	9 50	16 0			90		" "	7
65	1842	" ... 9	10 0	12 0			133	79	" "	7
66	1842	" ... 10	10 10	11 0			89		Sword handle of Perseus.....	7
67	1842	Nov. ... 13	15 0	16 0			46		" "	2
68	1844	Aug. ... 9	11 20	15 0	13 30		367	139	" "	4
69	1844	" ... 10	8 0	13 0			117		β Cassiopeæ	1
70	1844	" ... 11	9 50	15 0	14 0		622	152	" "	4
71	1844	" ... 12	13 50	15 0			46		" "	2
72	1845	" ... 9	10 0	13 0			81		Sword handle of Perseus.....	1
73	1846	" ... 10	12 0	14 0			46		" "	4
74	1846	" ... 11	9 0	10 0			41		" "	2
75	1847	" ... 10	10 0	12 0			41		Head of Perseus.....	1
76	1847	" ... 10	12 0	14 0			415		" "	4
77	1847	" ... 11	9 15	10 0			37		" "	4
78	1848	" ... 9	13 0	15 30	15 15		475	281	In Perseus	4
79	1848	" ... 9	13 30	15 30			216		" "	3
80	1848	" ... 9	15 0	16 45			55		In Perseus	1
81	1849	April... 19	13 0	14 0			54		" "	3
82	1849	Aug. ... 10	10 0	12 30			260		" "	4
83	1850	" ... 9	12 40	15 0	13 30		451	216	In Perseus.....	3
84	1850	" ... 10	10 0	12 0			312		" "	3
85	1850	" ... 10	12 0	15 0			351		" "	2
86	1852	" ... 9	14 0	14 40			19		" "	1
87	1853	" ... 10	11 45	15 25	14 30		408	119	Sword handle of Perseus.....	3
88	1855	" ... 9	10 25	14 30	13 30		385	110	" "	3
89	1855	" ... 10	10 30	13 0	13 30		290	136	" "	3
90	1855	" ... 11	10 0	11 0			37		" "	1
91	1855	" ... 11	13 55	15 5			58		" "	1
92	1855	" ... 12	13 30	14 45			36		" "	1
93	1856	" ... 10	11 5	14 50	12 30		283	100	" "	3
94	1858	" ... 9	10 45	15 10	14 35		128		" "	1
95	1858	" ... 9	12 5	14 5			56		" "	1
96	1858	" ... 10	8 30	9 0			11		" "	1
97	1858	" ... 10	13 7	13 52			29		" "	1
98	1858	" ... 11	14 15	15 15			33		" "	1
99	1859	July ... 29	10 30	11 30			16		" "	1
100	1859	Aug. ... 5	11 0	12 0			19		" "	1
101	1859	" ... 9	11 0	13 0			64		" "	3
102	1859	" ... 9	13 0	15 30	15 15		304	156	Sword handle of Perseus.....	4
103	1859	" ... 9	14 15	15 30			56		$\alpha = 38^\circ 30', \delta = +57^\circ 15'$	1
104	1860	" ... 9	10 0	15 10	14 30		588	155	Sword handle of Perseus.....	7

51. A. J. S. XXXVII, 325.
 52. Trans. Am. Phil. Soc. VII, 268.
 53. " " " " " "
 54. A. J. S. XXXVII, 325.
 55. Trans. Am. Phil. Soc. VII, 270.
 56. A. J. S. XXXIX, 328.
 57. " " " "
 58. " " " "
 59. A. J. S. XLII, 397.
 60. " " " "
 61. A. J. S. XLI, 399.
 62. " " " "
 63. A. J. S. XLIII, 212.
 64. A. J. S. XLIII, 377.

65. A. J. S. XLIII, 377.
 66. " " " "
 67. A. J. S. XLIV, 209.
 68. A. J. S. XLVIII, 316.
 69. A. J. S. XLVIII, 320.
 70. A. J. S. XLVIII, 316.
 71. " " " "
 72. A. J. S. I, 86.
 73. A. J. S. II, 125.
 74. " " " "
 75. Sidereal Messenger, II, 14.
 76. A. J. S. VI, 278.
 77. " " " "
 78. A. J. S. VI, 279.

Meteor Showers—Cont'd.

Ref. number.	Authority.	Place of observation.	Remarks.
51	E. C. Herrick.....	New Haven, Conn.	
52	C. G. Forshey	St. Louis, Mo.	
53	C. G. Forshey	Illinois River.	
54	E. C. Herrick.....	New Haven, Conn.	
55	C. G. Forshey	Philadelphia, Pa.	
56	E. C. Herrick.....	New Haven, Conn.....	Moon set at 14h. 0m.
57	G. C. Schaeffer	Jamaica, L. I.	
58	G. C. Schaeffer.....	Jamaica, L. I.	
59	C. G. Forshey	Vidalia, La.....	No trains; paths short.
60	E. C. Herrick	New Haven, Conn.	
61	Dr. J. S. Huntington, U. S. N.	Pensacola, Fla.	
62	Dr. John Locke	Cincinnati, Ohio.	
63	E. C. Herrick.....	New Haven, Conn.....	Moon set at 15h. 0m.
64	E. C. Herrick.....	New Haven, Conn.....	Cloudy; actual observing time 1h. 10m.
65	E. C. Herrick.....	New Haven, Conn.....	Cloudy.
66	E. C. Herrick.....	New Haven, Conn.....	Cloudy.
67	E. C. Herrick.....	New Haven, Conn.....	Cloudy after 16h.
68	E. C. Herrick.....	New Haven, Conn.....	Partially cloudy.
69	S. R. Williams.....	Canonsburg, Pa.	
70	E. C. Herrick.....	New Haven, Conn.	
71	E. C. Herrick.....	New Haven, Conn.	
72	E. C. Herrick.....	New Haven, Conn.....	Cloudy.
73	E. C. Herrick.....	New Haven, Conn.	
74	E. C. Herrick.....	New Haven, Conn.	
75	C. G. Forshey	Pass Christian, La.....	33 conformable with the radiant.
76	W. M. Smith	Manlius, N. Y	Cloudy after 14h.
77	E. C. Herrick.....	New Haven, Conn.....	Completely cloudy after 10h.
78	E. C. Herrick.....	New Haven, Conn.	
79	E. C. Herrick.....	"On Mt. Carmel."	
80	C. G. Forshey	Mouth of Miss. river.	
81	E. C. Herrick.....	New Haven, Conn.	
82	S. R. Williams.....	Canonsburg, Pa.	
83	E. C. Herrick.....	New Haven, Conn.	
84	E. C. Herrick.....	New Haven, Conn.	
85	E. C. Herrick.....	New Haven, Conn.	
86	John Edmunds.....	New Haven, Conn.	
87	E. C. Herrick.....	New Haven, Conn.	
88	E. C. Herrick.....	New Haven, Conn.....	306 conformable.
89	E. C. Herrick.....	New Haven, Conn.....	242 conformable.
90	E. C. Herrick.....	New Haven, Conn.....	One-half conformable.
91	E. C. Herrick.....	New Haven, Conn.....	45 conformable.
92	E. C. Herrick.....	New Haven, Conn.....	20 conformable.
93	E. C. Herrick.....	New Haven, Conn.	
94	F. Bradley	Observed on railroad between Daven-
95	Prof. A. C. Twining ...	Cleveland, Ohio.	port and Chicago, Ill.
96	Prof. A. C. Twining ...	Cleveland, Ohio.	
97	Prof. A. C. Twining ...	Cleveland, Ohio.	
98	Prof. A. C. Twining ...	Cleveland, Ohio.	
99	F. Bradley	Chicago, Ill	6 conformable to the August radiant.
100	F. Bradley	Chicago, Ill	8 conformable to the August radiant.
101	F. Bradley	Chicago, Ill.	
102	F. Bradley	Chicago, Ill	Only a few unconformable meteors.
103	Prof. A. C. Twining ...	Boston, Mass.	
104	E. C. Herrick.....	New Haven, Conn.	

79. A. J. S. VI₂, 279.
 80. A. J. S. XI₂, 133.
 81. A. J. S. VII₁₂, 429.
 82. " " "
 83. A. J. S. XI₂, 130.
 84. " " "
 85. " " "
 86. A. J. S. XIV₂, 430.
 87. A. J. S. XVI₂, 288.
 88. A. J. S. XX₂, 285.
 89. " " "
 90. " " "
 91. A. J. S. XX₂, 285.

92. A. J. S. XX₂, 285.
 93. A. J. S. XXII₂, 290.
 94. A. J. S. XXVI₂, 435.
 95. " " "
 96. " " "
 97. " " "
 98. " " "
 99. A. J. S. XXVIII₂, 446.
 100. " " "
 101. " " "
 102. " " "
 103. " " "
 104. A. J. S. XXX₂, 296.

Ref. number.	DATE.			TIME OF SHOWER.		Time of max. flight.	Whole number counted.	Max. hourly rate.	Radiant point.	No. of observ's.
	Year.	Month.	Day.	Beginning.	End.					
				<i>h. m.</i>	<i>h. m.</i>	<i>h. m.</i>				
105	1860	Aug.	9	11 30	13 0	57	Sword handle of Perseus.....	1
106	1860	"	10	11 0	14 0	12 30	381	146	$\alpha = 32^\circ, \delta = +61^\circ$	5
107	1860	Nov.	7	7 0	9 0	46	N. E. of zenith	3
108	1860	"	12	6 30	17 0	14 30	423	68
109	1860	"	12	10 0	16 0	14 30	381	90	In Leo.....	5
110	1860	"	13	6 15	17 0	13 30	500	83
111	1860	"	13	15 15	16 15	21	In the "Sickle" in Leo	1
112	1860	"	14	15 40	16 25	15	" " "	1
113	1860	Dec.	12	8 20	12 0	11 30	180	60
114	1861	April...	19	14 45	16 0	52	2
115	1861	July ...	3	9 5	10 5	36	4
116	1861	Aug. ...	10	8 0	15 15	13 30	397	75	δ Persei.....	2
117	1861	"	10	10 0	13 0	12 30	289	119	Sword handle of Perseus.....	4
118	1861	"	10	10 25	14 45	12 30	289	85	Between η , γ , and τ Persei ...	2
119	1861	"	10	10 30	13 0	95	In Perseus	1
120	1861	"	10	12 25	13 0	12 30	160	37	1
121	1861	"	11	8 15	10 15	52	2
122	1861	"	11	9 30	12 0	47	In Perseus	1
123	1861	Nov. ...	10	13 0	15 0	11	$\alpha = 156^\circ 30', \delta = +40^\circ 40'$	1
124	1861	"	11	14 15	16 40	32	Near γ Leonis.....	1
125	1861	"	11	14 15	15 30	32	In Leo.....	2
126	1861	"	11	16 0	17 0	15	"	1
127	1861	"	12	15 0	17 0	16 30	139	72	"	4
128	1861	"	12	15 0	16 30	15	1
129	1861	"	12	11 15	12 15	11	4
130	1861	"	12	15 20	16 20	27	4
131	1861	"	13	10 0	13 0	15	Zenith.....	1
132	1861	"	13	15 0	17 30	19	1
133	1861	"	13	15 15	17 38	23	In Perseus	1
134	1862	Aug. ...	9	14 0	15 40	90	In Perseus	2
135	1862	"	10	12 30	15 30	51	"	1
136	1862	Nov.	13	12 0	15 0	59	In Leo.....	2
137	1862	"	13	15 15	17 5	31	"	1
138	1862	"	13	16 30	17 30	17	1
139	1863	Aug. ...	10	8 30	10 0	41	1
140	1863	"	10	9 0	12 0	399	In Perseus.....	4
141	1863	"	10	9 0	13 0	257	2
142	1863	"	10	9 15	10 15	96	3
143	1863	"	10	10 0	13 50	130	1
144	1863	"	10	12 0	14 0	289	3
145	1863	"	10	13 0	14 0	87	1
146	1863	"	10	15 10	15 40	153	6
147	1863	"	11	9 0	11 0	67	1
148	1863	Nov.	11	10 0	15 0	105
149	1863	"	11	11 22	14 52	185
150	1863	"	12	10 0	14 0	129
151	1863	"	12	10 20	15 51	199
152	1863	"	13	8 0	14 0	107	8
153	1863	"	13	9 30	13 30	32
154	1863	"	13	10 10	17 7	15 0	213	46	In Leo.....	6
155	1863	"	13	10 38	17 16	15 30	316	69	In the "Sickle" in Leo.....	7

105. A. J. S. XXXI₂, 136.
 106. " " "
 107. A. J. S. XXXI₂, 137.
 108. A. J. S. XXXI₂, 138.
 109. A. J. S. XXXI₂, 139.
 110. A. J. S. XXXI₂, 138.
 111. A. J. S. XXXI₂, 137.
 112. " " "
 113. A. J. S. XXXI₂, 138.
 114. A. J. S. XXXI₂, 294.
 115. A. J. S. XXXI₂, 296.
 116. A. J. S. XXXI₂, 294.
 117. " " "

118. A. J. S. XXXII₂, 294.
 119. A. J. S. XXXII₂, 447.
 120. A. J. S. XXXII₂, 295.
 121. A. J. S. XXXIII₂, 148.
 122. A. J. S. XXXII₂, 447.
 123. A. J. S. XXXIII₂, 148.
 124. " " "
 125. A. J. S. XXXIII₂, 146.
 126. " " "
 127. " " "
 128. A. J. S. XXXIII₂, 147.
 129. A. J. S. XXXIII₂, 148.
 130. " " "

Meteor Showers—Cont'd.

Ref. number.	Authority.	Place of observation.	Remarks.
105	F. Bradley	Chicago, Ill.	
106	F. Bradley	Chicago, Ill.	
107	Prof. C. U. Shepard.....	Off Cape Hatteras.	
108	Francis Miller	Montgomery Co., Md.....	Several students assisted in the ob-
			servations.
109	Prof. D. Kirkwood	Bloomington, Ind.....	Half the number conformable after
			13h.
110	Francis Miller	Montgomery Co., Md.....	Several students assisted in the ob-
111	Prof. H. A. Newton.....	New Haven, Conn.	servations.
112	Prof. H. A. Newton.....	New Haven, Conn.	
113	Francis Miller	Montgomery Co., Md.....	Several students assisted in the ob-
			servations.
114	E. C. Herrick.....	New Haven, Conn.	No decided radiant.
115	E. C. Herrick.....	New Haven, Conn.	
116	F. W. Russell	Natick, Mass.	
117	E. C. Herrick.....	New Haven, Conn.....	A very large meteor at 11h. 30m.
118	B. V. Marsh	Burlington, N. J.....	A very large meteor at 11h. 23m.
119	Prof. A. C. Twining ...	New Haven, Conn.	
120	E. M. Gummere	Burlington, N. J.	
121	John Roberts	Madison, Ind.	
122	Prof. A. C. Twining ...	New Haven, Conn.	
123	F. W. Russell	Natick, Mass.	
124	F. W. Russell	Natick, Mass.	
125	Prof. A. C. Twining ...	New Haven, Conn.	
126	E. C. Herrick.....	New Haven, Conn.	
127	Prof. A. C. Twining ...	New Haven, Conn.	
128	S. J. Gummere	Burlington, N. J.	
129	Prof. D. Kirkwood	Bloomington, Ind.	
130	Prof. D. Kirkwood	Bloomington, Ind.	
131	F. W. Russell	Natick, Mass.	
132	S. J. Gummere	Burlington, N. J.	
133	B. V. Marsh	Germantown, Pa.....	Thirteen left trains.
134	F. W. Russell.....	Winchendon, Mass.	
135	B. V. Marsh	Germantown, Pa.	
136	Prof. S. J. Gummere...	Haverford College, Pa.	
137	B. V. Marsh	Germantown, Pa.	
138	Prof. A. C. Twining.....	New Haven, Conn.	
139	W. G. Bryant.....	Winchendon, Mass.	
140	F. W. Russell.....	Natick, Mass.	
141	J. G. Pinkham.....	Manchester, Me.	
142	Prof. H. A. Newton.....	New Haven, Conn.	
143	B. V. Marsh.....	Philadelphia, Pa.	
144	F. W. Russell.....	Natick, Mass.	
145	F. Bradley.....	Chicago, Ill.	
146	Prof. H. A. Newton.....	New Haven, Conn.	
147	F. W. Russell.....	Natick, Mass.	
148	Prof. O. N. Stoddard...	Oxford, Ohio.	
149	Prof. H. L. Smith.....	Kenyon College, Ohio.	
150	Prof. O. N. Stoddard...	Oxford, Ohio.	
151	Prof. H. L. Smith.....	Kenyon College, Ohio.	
152	Prof. A. D. Bache.....	Coast Survey Office, Wash-	More than half the tracks plotted.
		ington, D. C.	
153	Prof. H. A. Newton.....	New Haven, Conn.....	Hazy.
154	Capt. J. M. Gilliss,	U. S. Naval Observatory.	Track of each meteor plotted.
	U. S. N.		
155	Prof. S. J. Gummere...	Haverford College, Pa.....	200 tracks plotted.

131. A. J. S. XXXIII₂, 148.
 132. A. J. S. XXXIII₂, 147.
 133. " " "
 134. A. J. S. XXXIV₂, 295.
 135. " " "
 136. A. J. S. XXXV₂, 146.
 137. " " "
 138. " " "
 139. A. J. S. XXXVI₂, 306.
 140. A. J. S. XXXVI₂, 305.
 141. A. J. S. XXXVI₂, 304.
 142. A. J. S. XXXVI₂, 302.
 143. A. J. S. XXXVI₂, 304.

144. A. J. S. XXXVI₂, 305.
 145. " " "
 146. A. J. S. XXXVI₂, 302.
 147. A. J. S. XXXVI₂, 305.
 148. A. J. S. XXXVII₂, 144.
 149. A. J. S. XXXVII₂, 146.
 150. A. J. S. XXXVII₂, 144.
 151. A. J. S. XXXVII₂, 146.
 152. A. J. S. XXXVII₂, 145.
 153. A. J. S. XXXVII₂, 143.
 154. A. J. S. XXXVII₂, 141.
 155. A. J. S. XXXVII₂, 142.

CATALOGUE IV.—

Ref. number.	DATE.			TIME OF SHOWER.		Time of max. flight.	Whole number counted.	Max. hourly rate.	Radiant point.	No. of observations.
	Year.	Month.	Day.	Beginning.	End.					
				h. m.	h. m.	h. m.				
156	1863	Nov. ...	13	13 0	17 20	15 30	97	26	In the "Sickle" in Leo.....	2
157	1863	" ...	13	15 0	16 30		79		" " " " " " " " " " " "	1
158	1863	" ...	13	15 45	17 45		49		" " " " " " " " " " " "	1
159	1864	Aug. ...	9	9 0	16 0		17		" " " " " " " " " " " "	1
159 ₁	1864	" ...	9	10 0	11 0		33		" " " " " " " " " " " "	1
160	1864	" ...	9	10 20	13 0		29		" " " " " " " " " " " "	2
161	1864	" ...	9	10 30	13 0		332		" " " " " " " " " " " "	4
162	1864	" ...	9	10 30	13 0		300		" " " " " " " " " " " "	1
163	1864	" ...	9	10 40	12 0		13		" " " " " " " " " " " "	1
163 ₁	1864	" ...	9	11 30	14 30		44		" " " " " " " " " " " "	2
164	1864	" ...	9	12 0	12 30		25		" " " " " " " " " " " "	1
165	1864	" ...	9	13 0	15 30	14 30	691		" " " " " " " " " " " "	3
166	1864	" ...	9	15 0	16 0		50		" " " " " " " " " " " "	1
167	1864	Nov. ...	11	13 15	16 15		18		In Leo.....	1
168	1864	" ...	12	13 45	16 0		46		" " " " " " " " " " " "	1
169	1865	Aug. ...	9	14 25	15 50		19		$\alpha = 42^\circ, \delta = +57^\circ$	1
170	1865	" ...	12	10 45	12 45		16		$\alpha = 52^\circ, \delta = +58^\circ$	1
171	1865	" ...	15	11 5	15 5	14 30	178	61	" " " " " " " " " " " "	6
172	1865	Nov. ...	12	13 0	14 0		77		In the "Sickle" in Leo.....	1
173	1865	" ...	12	15 0	16 0		63		" " " " " " " " " " " "	1
174	1866	Aug. ...	9	9 0	9 20		14		" " " " " " " " " " " "	1
175	1866	" ...	9	12 0	14 0	12 30	114	76	" " " " " " " " " " " "	4
176	1866	" ...	10	9 0	10 15		50		In Cassiopeæ.....	1
177	1866	" ...	10	9 10	15 0	14 30	364	97	In Perseus.....	1
178	1866	" ...	10	9 15	11 30		100		" " " " " " " " " " " "	1
179	1866	" ...	10	12 0	14 15		129		" " " " " " " " " " " "	1
180	1866	" ...	10	12 0	14 15		164		" " " " " " " " " " " "	1
181	1866	" ...	10	13 0	14 0		51		" " " " " " " " " " " "	1
182	1866	" ...	10	14 0	16 0	14 30	75	43	In Perseus.....	1
183	1866	" ...	11	8 15	9 0		8		" " " " " " " " " " " "	1
184	1866	" ...	11	14 0	16 0		63		" " " " " " " " " " " "	1
185	1866	Nov. ...	8	9 45	13 20		93		"About Capella".....	1
186	1866	" ...	9	10 0	15 0		47		In Leo.....	3
187	1866	" ...	11	13 30	14 30		27		" " " " " " " " " " " "	1
188	1866	" ...	12	7 0	18 0	14 30	205	45	" " " " " " " " " " " "	1
189	1866	" ...	12	10 45	14 30	13 30	85	31	" " " " " " " " " " " "	4
190	1866	" ...	12	11 0	16 40	16 10	402	99	Between α and γ Leonis.....	1
191	1866	" ...	12	11 10	13 40	11 30	236	108	" " " " " " " " " " " "	12
192	1866	" ...	12	11 30	13 0		65		" " " " " " " " " " " "	5
193	1866	" ...	12	12 0	17 0	15 30	603	155	" " " " " " " " " " " "	5
194	1866	" ...	12	13 0	15 0		35		" " " " " " " " " " " "	3
195	1866	" ...	12	13 10	13 40		8		" " " " " " " " " " " "	1
196	1866	" ...	12	14 20	15 20		56		" " " " " " " " " " " "	3
197	1866	" ...	12	14 40	17 30	17 15	458	180	$\alpha = 147^\circ 30', \delta = +23^\circ 15'$	12
198	1866	" ...	12	15 54	16 39		64		" " " " " " " " " " " "	1
199	1866	" ...	12	16 6	16 36		8		" " " " " " " " " " " "	1
200	1866	" ...	12				354		" " " " " " " " " " " "	7
201	1866	" ...	13	7 0	18 0	13 30	453	71	" " " " " " " " " " " "	1
202	1866	" ...	13	8 30	16 30	15 0	440	138	" " " " " " " " " " " "	4
203	1866	" ...	13	10 0	15 0		150		" " " " " " " " " " " "	1
204	1866	" ...	13	11 0	14 0	13 0	492	202	" " " " " " " " " " " "	10
205	1866	" ...	13	11 0	16 10	15 30	901	212	In Leo.....	12
206	1866	" ...	13	11 20	14 40	12 30	261	113	" " " " " " " " " " " "	5

- | | | |
|--------------------|---------------------------------|------|
| 156. | A. J. S. XXXVII ₂ , | 142. |
| 157. | A. J. S. XXXVII ₂ , | 143. |
| 158. | " | " |
| 159. | A. J. S. XXXVIII ₂ , | 432 |
| 159 ₁ . | " | " |
| 160. | " | " |
| 161. | " | " |
| 162. | " | " |
| 163. | " | " |
| 163 ₁ . | " | " |
| 164. | " | " |
| 165. | " | " |
| 166. | " | " |
| 167. | A. J. S. XXXIX ₂ , | 229. |

168. A. J. S. XXXIX₂, 229.
169. A. J. S. XL₂, 284.
170. " " " "
171. " " " "
172. A. J. S. XLI₂, 273.
173. " " " "
174. A. J. S. XLII₂, 286.
175. A. J. S. XLII₂, 429.
176. A. J. S. XLII₂, 286.
177. A. J. S. XLII₂, 429.
178. " " " "
179. " " " "
180. " " " "
181. A. J. S. XLII₂, 429.

Meteor Showers—Cont'd.

Ref. number.	Authority.	Place of observation.	Remarks.
156	B. V. Marsh.....	Germantown, Pa.....	Hazy. { *On Nov. 13, 55 tracks were plotted in Phila. by H. D. Vail, 56 tracks were plotted in West Chester by B. Hooper, 27 tracks were plotted in Easton by E. Menline, 23 tracks were plotted in St. Louis by Prof. W. Chauvenet.
157	C. E. Dutton.....	Norfolk, Va.....	
158	J. H. Worrall.....	West Chester, Pa.....	
159	H. P. Tuttle.....	Off Charleston, S. C.....	
159 ₁	G. Scarborough.....	Riverside, Kan.....	
160	H. S. Osborn.....	Belvidere, N. J.....	
161	Francis Bradley.....	Chicago, Ill.....	Cloudy in the Middle and N. E. States.
162	W. H. R. Lykins.....	Lawrence, Kan.....	
163	B. V. Marsh.....	Philadelphia, Pa.....	Some left trains lasting 6 or 7 seconds. Off Martha's Vineyard, Mass. Williams College. No observations from 12h. 45m. to 13h. 0m. F. W. Russell, of Cambridge, Mass., observed several evenings in first half of November, and counted, in all, 875 meteors. Observed seven hours. Williams College.
163 ₁	Prof. H. A. Newton.....	New Haven, Conn.....	
164	G. Scarborough.....	Riverside, Kan.....	
165	Francis Bradley.....	Chicago, Ill.....	
166	G. Scarborough.....	Riverside, Kan.....	
167	R. H. Stretch.....	Virginia City, Nevada.....	
168	R. H. Stretch.....	Virginia City, Nevada.....	
169	Prof. A. C. Twining.....	Hinsdale, Mass.....	
170	Prof. A. C. Twining.....	Hinsdale, Mass.....	
171	Prof. H. A. Newton.....	New Haven, Conn.....	
172	O. B. Wheeler.....	Detroit, Mich.....	
173	O. B. Wheeler.....	Detroit, Mich.....	
174	D. Trowbridge.....	Hector, N. Y.....	
175	Prof. H. A. Newton.....	Sherburne, N. Y.....	
176	D. Trowbridge.....	Hector, N. Y.....	
177	F. W. Russell.....	Natick, Mass.....	
178	Isaac Pierson.....	Vineyard Sound.....	
179	B. V. Marsh.....	Germantown, Pa.....	
180	R. M. Gummere.....	Germantown, Pa.....	
181	Prof. H. A. Newton.....	Sherburne, N. Y.....	
182	J. H. Worrall.....	West Chester, Pa.....	
183	D. Trowbridge.....	Hector, N. Y.....	
184	J. H. Worrall.....	West Chester, Pa.....	
185	James Ferguson.....	U. S. Naval Observatory.....	
186	James Ferguson.....	U. S. Naval Observatory.....	
187	F. Bradley.....	Chicago, Ill.....	
188	Prof. Hopkins.....	Williamstown, Mass.....	
189	C. G. Rockwood.....	Newark, N. J.....	
190	James Ferguson.....	U. S. Naval Observatory.....	
191	Prof. H. A. Newton.....	New Haven, Conn.....	
192	F. Bradley.....	Chicago, Ill.....	
193	C. S. Lyman.....	New Haven, Conn.....	
194	Prof. Hinrichs.....	Iowa City, Iowa.....	
195	B. V. Marsh.....	Germantown, Pa.....	
196	O. B. Wheeler.....	Detroit, Mich.....	
197	Prof. H. A. Newton.....	New Haven, Conn.....	
198	Prof. D. Kirkwood.....	Canonsburg, Pa.....	
199	B. V. Marsh.....	Germantown, Pa.....	
200	Maria Mitchell.....	Vassar College, N. Y.....	
201	Prof. Hopkins.....	Williamstown, Mass.....	
202	Prof. Hinrichs.....	Iowa City, Iowa.....	
203	John T. Wheeler.....	Concord, N. H.....	
204	Prof. C. S. Lyman.....	New Haven, Conn.....	
205	Prof. H. A. Newton.....	New Haven, Conn.....	
206	C. G. Rockwood.....	Newark, N. J.....	

182. A. J. S. XLII₂, 429.
 183. A. J. S. XLII₂, 286.
 184. A. J. S. XLII₂, 429.
 185. A. J. S. XLIII₂, 78; Nov. Meteors, 1866,
 U. S. N. Obs'y, 8vo.
 186. " " "
 187. A. J. S. XLIII₂, 78.
 188. " " "
 189. " " "
 190. A. J. S. XLIII₂, 78, Nov. Meteors, 1866,
 U. S. N. Obs'y, 8vo.
 191. A. J. S. XLIII₂, 78.
 192. " " "
 193. " " "

194. A. J. S. XLIII₂, 78.
 195. " " "
 196. " " "
 197. " " "
 198. " " "
 199. " " "
 200. " " "
 201. " " "
 202. " " "
 203. Smithsonian Archives (not yet printed).
 204. A. J. S. XLIII₂, 78.
 205. " " "
 206. " " "
 *A. J. S. XXXVII₂, 141.

CATALOGUE IV.—

Ref. number.	DATE.			TIME OF SHOWER.		Time of max. flight.	Whole number counted	Max. hourly rate.	Radiant point.	No. of observ'rs.
	Year.	Month.	Day.	Begin-ning.	End.					
				<i>h. m.</i>	<i>h. m.</i>	<i>h. m.</i>				
207	1866	Nov.	13	12 0	12 36		28			1
208	1866	"	13	12 0	13 0		35		$\alpha = 147^{\circ} 30', \delta = + 24^{\circ} 30'$	1
209	1866	"	13	12 0	17 15	14 30	101	51		3
210	1866	"	13	14 0	16 0		180			4
211	1866	"	13	14 0	16 30	14 30	173	91	α Leonis.....	1
212	1866	"	13	15 8	16 8		43		Bend of the Sickle.....	6
213	1866	"	13				419			1
214	1866	"	16	13 0	15 0		24		Between μ and ϵ Leonis.....	1
215	1866	"	17	14 0	17 30		75		Near α Leonis.....	1
216	1867	May.	20				20		A little southeast of Lyra.....	1
217	1867	Aug.	8	12 15	12 55		17		In Perseus.....	1
218	1867	"	9	13 3	14 30		35		"	1
219	1867	"	9	13 3	14 30		27		"	1
220	1867	"	9	14 15	15 25		39		"	1
221	1867	"	10	14 0	15 40		44		"	1
222	1867	"	10	15 0	15 30		17		"	1
223	1867	"	11	14 0	15 30		45		"	1
224	1867	Nov.	13	9 0	17 0	15 30	535	276		5
225	1867	"	13	9 28	13 0		50			10
226	1867	"	13	10 0	16 0		4,937			
227	1867	"	13	11 0	16 0		1,000			4
228	1867	"	13	11 45	18 25	15 30	2,325	1,488		
229	1867	"	13	12 0	16 40	15 45	2,056			
230	1867	"	13	12 0	17 45	15 50	3,730	2,184		2
231	1867	"	13	12 30	17 0	15 52	1,713	1,076	$\alpha = 147^{\circ} 45', \delta = + 23^{\circ} 0'$	
232	1867	"	13	13 0	18 0	16 30	3,044	2,110	γ Leonis.....	15
233	1867	"	13	13 0	13 30		500			4
234	1867	"	13	13 0	16 28	16 22	1,474	2,220	In Leo.....	1
235	1867	"	13	13 0	18 0	16 10	2,267	1,472		4
236	1867	"	13	14 0	18 13	16 30	1,626	1,082		1
237	1867	"	13	15 3	17 45		1,800		In Leo.....	4
238	1867	"	13	15 45	16 45		1,088		$\alpha = 150^{\circ} 45', \delta = + 21^{\circ} 55'$	2
239	1867	"	13	15 55	16 35		339		$\alpha = 148^{\circ} 0', \delta = + 23^{\circ} 0'$	4
240	1867	"	13	16 0	17 0		500			1
241	1867	"	13	16 5	17 54		1,301		In Leo.....	1
242	1867	"	13	16 30	17 30		460			1
243	1867	"	13	17 0	17 22		100		In Leo.....	1
244	1867	"	13	17 0	17 30		61			1
245	1867	"	13				500			2
246	1868	Aug.	9	12 15	14 0		56		In Perseus.....	3
247	1868	Nov.	13	10 0	15 30		1,330			2
248	1868	"	13	10 45	18 0	16 30	2,886	572		1
249	1868	"	13	11 0	18 11		2,500			3
250	1868	"	13	11 30	17 30		1,462		$\alpha = 152^{\circ}, \delta = + 18^{\circ}$	10
251	1868	"	13	11 35	13 20		700			3
252	1868	"	13	12 0	17 54	17 27	5,573	1,402	In Leo.....	2
253	1868	"	13	11 34	17 40	17 0	5,000	1,400		1
254	1868	"	13	12 29	14 20		800			1
255	1868	"	13	12 51	17 15	17 10	5,670	1,545		1
256	1868	"	13	13 0			896			1
257	1868	"	13	13 0	17 28		850			4
258	1868	"	13	13 0	18 0	16 30	1,341	628		3
259	1868	"	13	13 4	17 37		1,926			

207. A. J. S. XLIII₂, 78.

208. " " " "

209. A. J. S. XLIII₂, 413.210. A. J. S. XLIII₂, 78.211. A. J. S. XLIII₂, 78; Nov. Meteors, 1866, U. S. N. Obs'y, 8vo.212. A. J. S. XLIII₂, 78.

213. " " " "

214. A. J. S. XLIII₂, 78; Nov. Meteors, 1866, U. S. N. Obs'y, 8vo.

215. " " " "

216. Trans. St. Louis Acad. Sci., II, 577.

217. A. J. S. XLIV₂, 426.

218. " " " "

219. " " " "

220. A. J. S. XLIV₂, 426.

221. " " " "

222. " " " "

223. " " " "

224. A. J. S. XLV₂, 78.

225. " " " "

226. A. J. S. XLV₂, 225.227. A. J. S. XLV₂, 78.

228. Smithsonian Archives (not yet printed).

229. A. J. S. XLV₂, 225.230. A. J. S. XLV₂, 78.

231. " " " "

232. Proc. Am. Phil. Soc. X, 356.

233. A. J. S. XLV₂, 78.

234. " " " "

Meteor Showers—Cont'd.

Ref. number.	Authority.	Place of observation.	Remarks.
207	B. V. Marsh	Germantown, Pa.	
208	Prof. A. C. Twining.....	New Haven, Conn.	
209	The Denver News.....	Denver, Col.	
210	W. A. Anthony.....	Franklin, N. Y.	
211	James Ferguson.....	U. S. Naval Observatory.	
212	Prof. A. C. Twining.....	New Haven, Conn.	
213	Maria Mitchell.....	Vassar College, N. Y.	
214	James Ferguson.....	U. S. Naval Observatory.....	120 tracks plotted on charts from Nov. 9 to Nov. 17, inclusive.
215	James Ferguson.....	U. S. Naval Observatory.	Observations made in the evening.
216	R. Hayes	St. Louis, Mo.....	"Moon two days past the full."
217	F. W. Russell.....	Winchendon, Mass.....	
218	B. V. Marsh	Philadelphia, Pa.	
219	C. H. Darlington.....	Philadelphia, Pa.	
220	Lewis Swift.....	Marathon, N. Y.	
221	F. W. Russell.....	Winchendon, Mass.	
222	Lewis Swift.....	Marathon, N. Y.	
223	F. W. Russell.....	Winchendon, Mass.	
224	Prof. T. C. Wylie.....	Bloomington, Ind.	
225	Prof. H. A. Newton	New Haven, Conn.	Cloudy from 12h. 0m. to 15h. 15m.
226	Prof. N. R. Leonard.....	Iowa City, Iowa.....	
227	S. J. Gummere	Haverford College, Pa.	
228	Prof. F. H. Snow.....	Lawrence, Kan.	
229	Prof. C. A. Young.....	Hanover, N. H.....	Dartmouth College.
230	Prof. T. H. Safford.....	Chicago, Ill.....	Number of observers varied from 8 to 30.
231	Francis Bradley.....	Evanston, Ill.	Several observers.
232	James McClure	Philadelphia, Pa.....	
233	San Francisco Times.....	San Francisco, Cal.	
234	Prof. H. A. Newton.....	New Haven, Conn.....	One-third of the sky covered with clouds.
235	G. F. Kingston.....	Toronto, Canada.	
236	Robert B. Taber.....	New Bedford, Mass.	
237	Profs. Newcomb and Eastman.	U. S. N. Observatory	147 tracks mapped.
238	Prof. J. C. Watson.....	Ann Arbor, Mich.	
239	Prof. W. Harkness.....	Richmond, Va.	
240	Mrs. J. H. Trumbull.....	Hartford, Conn.	
241	Prof. G. W. Hough.....	Dudley Observatory	*In Chihuahua, Mexico, the meteors fell so fast they could not be counted; often 20 or 30 visible at once.
242	J. N. Flint.....	Canaseraga, N. Y.	
243	Prof. A. C. Twining.....	New Haven, Conn.	
244	J. D. Parker.....	Topeka, Kan.	
245	Prof. E. Loomis	New Haven, Conn.....	Observed one hour.
246	C. G. Rockwood	Durham, Conn.	
247	Robert B. Taber.....	New Bedford, Mass	Actual observing time 4h. 35m.
248	G. T. Kingston.....	Toronto, Canada.	
249	T. A. Wylie.....	Bloomington, Ind.	
250	W. S. Gilman, Jr.....	Palisades, N. Y.....	†Several bright meteors were seen during the Nov. shower in 1868.
251	B. J. Gilman.....	Williamstown, Mass.	
252	Prof. H. A. Newton.....	New Haven, Conn.	
253	S. J. Gummere	Haverford College, Pa.	
254	Prof. J. R. Eastman	U. S. N. Observatory.	
255	C. G. Rockwood	Brunswick, Me.	
256	Lewis Swift.....	Marathon, N. Y.....	The time when observations ended not given.
257	P. E. Chase.....	Haverford College, Pa.	
258	W. H. Pratt.....	Davenport, Iowa	43 were counted before 13h. 0m.
259	C. G. Berner.....	Vevay, Ind.	

235. A. J. S. XLV₂, 78.
 236. " " "
 237. A. J. S. XLV₂, 225; Nov. Meteors, 1867, U. S. N. Obs'y, 8vo.
 238. A. J. S. XLV₂, 78.
 239. A. J. S. XLV₂, 225; Nov. Meteors, 1867, U. S. N. Obs'y, 8vo.
 240. A. J. S. XLV₂, 78.
 241. " " "
 242. " " "
 243. " " "
 244. " " "
 245. " " "
 246. A. J. S. XLVII₂, 287,
 247. A. J. S. XLVII₂, 118.
 248. " " "

249. A. J. S. XLVII₂, 118.
 250. " " "
 251. " " "
 252. " " "
 253. " " "
 254. Nov. Meteors, 1868, U. S. N. Obs'y, 8vo.
 255. A. J. S. XLVII₂, 118.
 256. " " "
 257. A. J. S. XLVII₂, 118; Proc. Am. Phil. Soc. X, 539.
 258. A. J. S. XLVII₂, 118; Dav. Acad. Nat. Sci. I, 14.
 259. A. J. S. XLVII₂, 118.
 * A. J. S. XLV₂, 78.
 † A. J. S. XLVII₂, 399.

CATALOGUE IV.—

Ref. number.	DATE.			TIME OF SHOWER.		Time of max. flight.	Whole number counted.	Max. hourly rate.	Radiant point.	No. of observations.
	Year	Month.	Day.	Begin-ning.	End.					
				<i>h. m.</i>	<i>h. m.</i>	<i>h. m.</i>				
260	1868	Nov	13	14 20	17 58	14 30	3,766	900		5
261	1868	"	13	15 0	17 0	16 55	4,278	1,650		4
262	1868	"	13	15 7	17 38		1,780			3
263	1868	"	13	15 30	18 0		455			1
264	1868	"	13	16 0	17 15		711			1
265	1868	"	13	16 30	17 30		250			1
266	1868	"	13	7 0	19 0		833			2
267	1869	"	13	13 18	15 43		830			4
268	1869	"	12	12 30	15 30		556			2
269	1870	"	13	11 5	15 45		31		In Leo.	3
270	1870	"	13	13 20	16 10		153		"	6
271	1870	"	13	13 30	15 0		82		"	3
272	1870	"	13	13 30	15 0		13			1
273	1870	"	13	13 30	15 0		11			1
274	1871	Aug.	10	11 40	13 18		283			6
275	1871	Nov.	13				98			
276	1872	Aug.	9	10 30	15 30		358			4
277	1872	"	9	14 45	15 55		62			1
278	1872	Nov.	24	7 30	12 30		217		2° or 3° N. of γ Andromedæ	2
279	1872	"	25	10 25	14 0		80			5
280	1872	"	27	6 5	10 0	6 30	720	300	μ Andromedæ	1
281	1872	"	27	6 10	11 45		143		$\alpha = 15^\circ, \delta = +30^\circ$	1
282	1872	"	27	6 38	8 48	6 55	1,750	1,300	$\alpha = 25^\circ, \delta = +43^\circ$	4
283	1875	Aug.	5	13 30	15 30		112		In Perseus	1
284	1876	Oct.	24	11 30	15 30		68		Bet. Orion and the Pleiades.	1
285	1876	Nov.	13	12 0	13 0		12			1
286	1877	"	13	13 55	15 45		54		In Leo.	2
287	1885	"	27	6 0	6 24		44			1
288	1885	"	27	9 0			12		γ Andromedæ	3
289	1885	"	27	10 15	10 30		13			1
290	1885	"	27	7 15	7 45		100		2° N. W. of γ Andromedæ	1
291	1885	"	27	6 30	7 50		213			3
292	1885	"	27	7 0	9 0		328			4

260. A. J. S. XLVII₂, 118.261. Nov. Meteors, 1868, U. S. N. Observatory,
8 vo.262. A. J. S. XLVII₂, 118.

263. " " "

264. " " "

265. " " "

266. " " "

267. A. J. S. XLIX₂, 244.268. A. J. S. XLIX₂, 244.269. A. J. S. I₃, 30.

270. " " "

271. " " "

272. " " "

273. " " "

274. A. J. S. II₃, 227.275. A. J. S. II₃, 470.276. A. J. S. IV₃, 244.

Meteor Showers—Cont'd.

Ref. number.	Authority.	Place of observation.	Remarks.
260	Maria Mitchell	Vassar College, N. Y.	
261	Prof. J. R. Eastman ...	U. S. N. Observatory.....	90 tracks plotted.
262	B. J. Gilman.....	Williamstown, Mass.	
263	C. W. Tuttle.....	Boston, Mass.....	No observations between 16h. 4m. and
264	Robert B. Taber.....	New Bedford, Mass.	16h. 33m.
265	J. E. Hendricks.....	Des Moines, Iowa.	
266	B. F. Mudge.....	Manhattan, Kan.	
267	Frederickton, N. B.....	Reported by Prof. H. A. Newton.
268	George Davidson.....	Santa Barbara, Cal.	
269	Prof. H. A. Newton.....	New Haven, Conn.	
270	Prof. H. A. Newton	New Haven, Conn.	
271	C. G. Rockwood	Brunswick, Me.....	6 tracks plotted.
272	B. V. Marsh.....	Burlington, N. J.	
273	J. G. Gummere	Burlington, N. J.	
274	Prof. H. A. Newton.....	Sherburne, N. Y.	
275	Prof. H. A. Newton.....	New Haven, Conn.....	Only a few belonged to the November
276	Prof. C. A. Lyman.....	New Haven, Conn.	system.
277	R. W. McFarland	Oxford, Ohio.	
278	Prof. H. A. Newton.....	New Haven, Conn.	
279	Prof. H. A. Newton.....	New Haven, Conn.	
280	Prof. J. R. Eastman ...	U. S. N. Observatory.	
281	B. V. Marsh.....	Philadelphia, Pa.	
282	Prof. H. A. Newton.....	New Haven, Conn.	
283	Prof. J. R. Eastman ...	U. S. N. Observatory.....	Most of the meteors had short trains.
284	Prof. J. R. Eastman ...	U. S. N. Observatory.....	A few left long trains.
285	Prof. H. A. Newton.....	New Haven, Conn	Night mostly cloudy; 3 conformable to radiant.
286	Prof. D. Kirkwood	Bloomington, Ind	Night mostly cloudy.
287	Robert Brown.....	New Haven, Conn.	
288	H. A. Newton.....	New Haven, Conn	Hourly number estimated for 3 ob- servers, 100.
289	H. A. Newton.....	New Haven, Conn.	
290	C. A. Young	Princeton, N. J.	
291	A. Hall	Georgetown, D. C.	
292	D. Horgan	Georgetown, D. C.	

277. A. J. S. IV₃, 244.

278. A. J. S. V₃, 53.

279. " " "

280. " " "

281. " " "

282. " " "

283. Astron'l and Meteor'l Observ'ns, 1875.

284. Astron'l and Meteor'l Observ'ns, 1876.

285. A. J. S. XII₃, 473.

286. A. J. S. XV₃, 76.

287. A. J. S. XXXI₃, 78.

288. " " "

289. " " "

290. " " "

291. " " "

292. " " "

CATALOGUE V.-Sporadic Meteors.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day.	Hour and minute.				
1	1779	Oct.	31	<i>h. m. s.</i> 6 10 0	Williamsburg, Va.	<i>s.</i>	Fell and disappeared below the horizon 4° N. of W. N. 20° E. Alt., 30°; az., N, 45° E.
2	1813	Mar ...	21	10 0 0	New Haven, Conn.	Like the moon	8 or 10	Alt., 35°; az., N. 20° E. Alt., 30°; az., N, 45° E.
3	1819	July ...	24	Youngstown, Ohio.	Alt., 45°; az., N. 45° E.
4	1819	Nov. ...	21	Evening ...	Easton, Penn.	Alt., 43°; az., S. 12° E. Alt., 23°; az., S.
5	1822	Mar. ...	9	10 0 0	New York, N. Y.	5	Alt., 60°; az., N. 30° E.
6	1823	Mar. ...	31	7 30 0	New Haven, Conn.	White tinged with blue.	2.5	Alt., 9° 48'; az., N. 41° 54' E. Alt., 3° 6'; az., N. 26° 57' E.
7	1826	Apr. ...	1	9 0 0	New Haven, Conn.	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.
8	1826	Apr. ...	14	11 20 0	Burlington, Vt.	At appearance 7' and at disappearance 28' in diameter.	2	Alt., 15° E. Alt., 40°; az., N.
9	1828	Feb.	11	11 12 0	On the East river, at New York.	Diameter 15'	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.
10	1836	Aug. ...	20	4 0 0	Eight miles S. W. of Jack- sonville, Ill.	Like a globe of fire.	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.
11	1837	July ...	5	7 8 0	Rochester, N. H.	Like "white-heated iron."	1m.	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.
12	1837	Dec. ...	14	7 39 32	New Haven, Conn.	1.0 or 1.5	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.
13	1838	May ...	18	8 0 0	Throughout the northern United States and part of upper Canada.	From 1/4 to 3/4 the size of the moon.	7	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.
14	1840	May ...	12	15 0 0	Four miles south of Ston- ington, Conn.	One-fourth the diameter of the moon.	3m. 30s.	Alt., 60°; az., N. 15° E. Alt., 40°; az., N.

1. Trans. Am. Phil. Soc. II (O. S.), 173, 175.
2. A. J. S. XII, 35.
3. A. J. S. VI, 315.
4. A. J. S. VI, 316.
5. A. J. S. VI, 318.
6. A. J. S. XI, 184.
7. A. J. S. " "
8. A. J. S. XI, 120.
9. A. J. S. XIV, 199.
10. A. J. S. XXXIII, 402.
11. A. J. S. XXXIII, 200.
12. A. J. S. XXXVII, 130.
13. A. J. S. XXXV, 223.
14. A. J. S. XXXIX, 382.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
1	Train visible 15m.....	0	Seen in Philadelphia by David Rittenhouse.....	John Page.
2	Train 12° long.....	Easterly nearly horizontal..	The light was powerful; cast deep shadows. In 8 or 10 minutes after its disappearance a heavy report, accompanied by a decided jar, was heard.	S. E. Dwight.
3	Moving towards the north..	Disappeared with an explosion, the report of which was heard in about three minutes. At Gustavus, Ohio, the same object was seen and the report heard in about one minute.	Dr. H. Manning.
4	Towards the S. or S.W.....	Disappeared at an altitude of 35° with a loud explosion, the report of which was heard in "two, three, or four minutes."	Samuel Turney.
5	This meteor was seen in Canada, throughout New England and New York, and in several places in Pennsylvania. At Troy, N. Y., the report of the explosion was heard in 7m. 30s.	
6	Train several degrees long.....	Report of explosion heard in 4m. 25s.	
7	Train visible one minute.....	Disappeared at the zenith...	Report heard in two minutes; also seen at Arlington, Vt., where it seemed to move from north to south, and vanished at an altitude of 30°. Report heard in 1m. 30s.	Dr. H. S. Waterhouse.
8	N. E. from a point 50° below the zenith."	No explosion.	Prof. B. D. Silliman.
9	Train of same color as the meteor.	20	Vertically towards the earth	The meteor seemed to burst into fragments, but no report was heard.	R. Gaylord.
10	Left a "cloud of smoky appearance."	20	Course towards the N. N. W. and inclined to the horizon.	Report of explosion at disappearance distinctly heard.....	
11	Left a greyish train which, after waving and expanding for several minutes, gradually vanished.		
12	Brilliant train, visible a long time.	Downward at an angle of from 30° to 50° with the vertical.	Seemed to explode, but no report was heard. Position at appearance not well observed.	E. C. Herrick.
13	A brilliant train broken into several sections.	Nearly north and horizontal	Data collected and computations made by Prof. E. Loomis...	Profs. Hubbard, Catlin, and Barrows, and Messrs. R. W. Haskins, E. Brown, and J. and L. Chandler.
14	Brilliant train lasting several seconds after the extinction of the meteor.	Downward at an angle of 30° with the horizon.	Exploded with a loud report, which was heard in ten or fifteen seconds.	J. J. Comstock.

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day.					
			Hour and minute.					
			<i>h. m.</i>					
15	1841	Nov. ...	10 8 0	Boston, Mass.....	Preceding portion white; the following blue.	s.	Passed several degrees west of Fomalhaut and disappeared at an altitude of 10°.
16	1841	Nov. ...	10 8 0	New Haven, Conn.....	Diameter 3' or 4'.....	White	2 or 3	Appeared 30' below the Pleiades and disappeared midway between β and θ Ceti.
17	1845	Aug. ...	11 6 5	Cambridge, Mass.....	Estimated diameter less than 5'.	White	1	Alt., 25° 30'.
18	1845	Aug. ...	25 8 0	Cambridge, Mass.....	Half the diameter of the full moon.	Red and blue.....	Alt., 45°; az., S. 10° W.
19	1845	Aug. ...	31 14 20	Fayetteville, N. C.....	"As large as the sun."	20	Alt., 10°; az., S. 69° W. Alt., 8°; az., N. 88° W.
20	1846	July ...	13 9 30	York and Lancaster, Pa...	From one-third to two-thirds the size of moon.	Red and yellowish white.
21	1846	July ...	20 9 55	Cambridge, Mass.....	"Equal to Venus".....	Blue, red, and white	2	Alt., 35°; az., N. 83° W. Alt., 30°; az., N. 63° W.
22	1846	Aug. ...	11 10 10	Cambridge, Mass.....	"Brighter than Venus"...	Alt., 30°.
23	1848	Jan.....	20 5 0	Rome, N. Y.....	Brilliant white	Alt., 31°; az., N. 12° E. Alt., 16°; az., N. 12° W.
24	1848	Mar. ...	5 14 30	Nantucket, Mass.....	20' in diameter	Exploded and disappeared in a 2h. 30m. and $\delta + 27^{\circ}$.
25	1848	April...	15	Whitesville, Miss.	"As large as Venus".....	Bluish.	From 60 to 90m.	Alt., 60°; az., S. of W.
26	1850	June..	16 6 25	New Haven, Conn.....	Orange	First seen at a point midway between the zenith and horizon; az., S. 30° or 35° W.
* 27	1850	Sept. ...	30 8 54	Boston, Mass.....
28	1850	Sept....	30 8 53	Springfield, Mass.....
29	1850	Oct. ...	3 8 30	Hartford, Conn.....
30	1856	July ...	8 6 0	Thornhill, Hancock Co., Ala.	"6 or 8 inches in diam'r"

15. A. J. S. XLIII, 309.
 16. " " " " " "
 17. Proc. Am. Acad. Arts and Sci. I, 21.
 18. " " " " " "
 19. A. J. S. XLIX, 408.
 20. A. J. S. XLIX, 347.
 21. Proc. Am. Acad. Arts and Sci. I, 22.
 22. " " " " " "
 23. A. J. S. V, 437.
 24. Proc. Am. Acad. Arts and Sci. I, 329.
 25. A. J. S. VI, 148.
 26. A. J. S. XI, 131.
 27. " " " " " "
 28. " " " " " "
 29. Proc. A. A. S., 1851, 191.
 30. A. J. S. XXII, 448; XXIII, 138, 237.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
15	Left a train 2° or 3° long near and a little above δ Aquarii, visible more than 90s.	50	Towards the S. W.	W. C. Bond and Son.
16	Train 8° or 10° long and visible 3m.	7	Same meteor as the preceding.	W. C. Bond.
17	10 or 20	Toward the horizon at an angle of 50°	W. C. Bond.
18	From N. E. to S. E.	Exploded with a very loud report, which was heard in five or six minutes.	D. M. Ettinger.
19	Train 1° long, seen only at York.	From 77 Cygni to near a Cassiopeæ	Seen also at New Haven and many other places.	W. C. Bond.
20	gasl.	W. C. Bond.
21	A cometary tail of dense white light.	From a Cassiopeæ to 80 Pe-	Light of train could be seen through light cirrus clouds.	Edward Huntington.
22	Brilliant train, a portion visible 12m. or 13m.	wards the E.	Loud report heard in 92s. after the meteor disappeared over the ocean.	William Mitchell.
23	Disappeared with a loud explosion. Light strong enough to cast deep shadows, even in the moonlight.	George Rice and J. P. Humaston.
24	Left a bright train	At the time of appearance the sun was about 15° above the horizon and unobserved.	W. C. Bond.
25	Train 1° long.	The train was examined with the Cambridge equatorial, and it presented the appearance of minute bright clouds resembling cirro-cumuli.	J. C. Hoadley.
26	Left a brilliant train extending from the head of Medusa to 3° below α Arietis, visible more than one hour.	Horizontally from E. to W.	Same meteor as the preceding
27	The train was yellowish, and remained stationary for about 4m. It then slowly changed its form and its direction of motion, and at 9h. 18m. it was barely visible as a faint white cloud in the lower portion of Arles.	Disappeared below S. E. horizon. No explosion. Train swept over the moon's disk.	Gaylord Wells, by Prof. Brocklesby.
28	Train 15 or 16 feet long	From W. to S. E.	Seen also at Columbus, Grenada, and Holly Springs, Miss., and at Marion, Ala.	T. M. Peters and B. R. Delgraffenried.
29	White train, visible 15m.	35	Nearly vertical
30

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year	Month.	Day.					
			h. m.					
31	1857	April...	11 8 50	Near Lake Winnibigoshish, Minn.	"Greater than the full moon."	8.	Appeared in Hydra and moved W. to within a short distance of the horizon.
32	1859	Aug. ...	11 7 20	From Newburyport, Mass., to Petersburg, Va.	"As large as the sun"	Red.	4	Exploded over the vicinity of Cape May.
33	1859	Nov. ...	14 21 30	From Lake Michigan to Nantucket, Mass., and from Maine to Virginia.	Nearly as large as the sun.	Like a ball of fire ..	2
34	1860	July....	20		Nearly as large as the moon.	Computed altitudes: Over Lake Michigan, 120 miles. " Lake Huron, 85 miles. " Buffalo, N. Y., 62 miles. " Elmira, N. Y., 51 miles. " Long Island Sound, 42 miles.
35	1860	Aug. ...	2 Bet'n 10 and 11h.	Knoxville, Tenn.	1 or 2	Appeared 1° S. W. of ε Cygni and passed near the Dolphin.
36	1860	Aug. ...	6 Bet. 7.30 and 8h.	New York and New Haven, Conn.	In the S. E. Over the Atlantic Ocean.
37	1861	Aug. ...	10 11 30	New Haven, Conn.
38	1861	Oct. ...	4 7 20	New Haven, Conn.
39	1861	Dec. ...	8 18 0	Brunswick, Me.
40	1861	Dec. ...	17 4 45	Buffalo, N. Y.
41	1861	Dec. ...	25 "Sunset"	Southern Connecticut.....
42	1861	Dec. ...	30 7 0	Hartford, Conn.
43	1862	Jan. ...	2 19 17	New York, N. Y.	One-third diam'r of moon.
44	1862	Jan. ...	5 "Sunset"	Setauket, N. Y.
45	1862	Aug. ...	13 7 47	U. S. Naval Observatory...	"Sparkling white."	3	Near Pleiades.
46	1862	Sept. ...	3 12 0	U. S. Naval Observatory...	4	In the N.
47	1862	Oct. ...	3 12 15	U. S. Naval Observatory...
48	1862	Nov. ...	3 7 0	U. S. Naval Observatory...
49	1862	Dec. ...	11 7 0	U. S. Naval Observatory.

31. A. J. S. XXIV^a, 158.
 32. A. J. S. XXXVII^a, 300.
 (A. J. S. XXXIX^a, 137, 298, 447.
 33. { Jour. Frank. inst. XXXIX^a, 205, 253.
 { A. J. S. XXX^a, 156.
 34. A. J. S. XXX^a, 293; Smithsonian Contribu-
 tions, XVI.
 35. A. J. S. XXX^a, 295.
 36. A. J. S. XXX^a, 296; XXXII^a, 344.
 37. A. J. S. XXXI^a, 448.
 38. A. J. S. XXXII^a, 443.
 39. A. J. S. XXXII^a, 291.
 40. " " "
 41. " " "

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
31	White train lasting 5m. or 10m.....	0	Nearly horizontal.....	B. F. Odell.
32	Seen in New York, Connecticut, and Vermont. At its disappearance the explosion jarred the buildings in Troy, N. Y. The sun was shining brightly.	
33	No luminous train was seen, but a trail of smoke was left in its path.	15 or 20	Nearly vertical.....	Position computed by Prof. E. Loomis.	
34	In the vicinity of Elmira, N. Y., the meteor separated by explosion into two parts, which continued their course until over Nantucket, when by another explosion they were separated into small fragments, which continued their course as before.	Positions computed by Prof. C. S. Lyman.
35	From E. to W.....	Seen throughout the southern portion of the United States..	W. C. Kane.
36	From S. to N.	Also seen at Burlington, N. J., by B. V. Marsh.....	Prof. A. C. Twining.
37	Brilliant train, visible more than 15s.	29	Slightly inclined towards the earth.	
38	Brilliant train reaching from Cygnus to Cassiopeia. It slowly assumed a serpentine shape and remained visible 2m.	From west to east, nearly through the zenith.	
39	From W. to E.....	Broke into three fragments.	
40	Left a short train.	From an altitude of 10° obliquely to horizon.	
41	From a Cygni to Delphinus.	Explored with a report like a pistol.	
42	From E. to W.	
43	From the N. E.	
44	From N. W. to S. E.	
45	From E. to W.	
46	From N. E. to N. W.	
47	Train visible 10s.	From zenith to S. W.	
48	
49	

42. A. J. S. XXXIII, 291.
 43. " " " "
 44. " " " "
 45. Wash'n Ast. & Met. Obs'ns, 1862.
 46. " " " "
 47. " " " "
 48. Wash'n Ast. & Met. Obs'ns, 1862.
 49. " " " "

CATALOGUE V.—Sporadic Meteors—Continued.

Rel. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day.					
			Hour and minute.					
			<i>h. m.</i>					
50	1862	Dec. ...	16	U. S. Naval Observatory...	2.	
51	1862	Dec. ...	18	U. S. Naval Observatory...	5	
52	1862	Dec. ...	20	U. S. Naval Observatory...	5	
53	1862	Dec. ...	20	U. S. Naval Observatory...		
54	1863	Jan. ...	11	U. S. Naval Observatory...	5	
55	1863	April... 2	12	U. S. Naval Observatory...	5	
56	1863	April... 19	7	Philadelphia, Pa.	3 to 6	
57	1864	July ...	8	U. S. Naval Observatory...		
58	1864	July ...	13	U. S. Naval Observatory...		Appeared just below Delphinus.
59	1864	Oct. ...	9	Boston, Mass.		Appeared 80° S. of Mars.
60	1864	Nov. ...	10	U. S. Naval Observatory...		Appeared 70° W. of the zenith.
61	1865	Jan. ...	8	U. S. Naval Observatory...		In the N. E.
62	1865	Feb. ...	21	U. S. Naval Observatory...		In the N.
63	1865	Nov. ...	25	U. S. Naval Observatory...		
64	1866	July ...	26	Hector, N. Y.		
65	1866	Nov. ...	5	Glen Cove, N. Y.		
66	1867	Mar. ...	29	U. S. Naval Observatory...		In Cygnus.
67	1867	April ...	7	U. S. Naval Observatory...		Seen to explode in N. N. E. at alt. of 10 or 14°.
68	1867	July ...	18	Salem, Pa.		In the W.
69	1867	Oct. ...	15	Haverford College, Pa.		Appeared in the S. E., disappeared in the N. E.
70	1868	Feb. ...	6	U. S. Naval Observatory...		30° above eastern horizon.
71	1868	June...	11	Manhattan, Kan.		Appeared 30° N. of Pleiades.
72	1869	Jan. ...	20	U. S. Naval Observatory...		Appeared in W. at alt. 55°, disappeared at alt. 12°.
73	1869	May ...	20	New York, N. Y.		at alt. 25°; az., N. 20° W.
74	1869	Aug. ...	24	U. S. Naval Observatory...		Appeared in E.; alt., 30°; disappeared in N. 60° E.
75	1869	Oct. ...	26	Patterson, Ky.		From Mizar to 10° below Polaris.

50. Wash'n Ast. & Met. Obs'ns, 1862.
 51. " " " "
 52. " " " "
 53. " " " "
 54. Wash'n Ast. & Met. Obs'ns, 1863.
 55. " " " "
 56. A. J. S. XXXVII, 154.
 57. Wash'n Ast. & Met. Obs'ns, 1864.
 58. A. J. S. XXXVIII, 295.
 59. Wash'n Ast. & Met. Obs'ns, 1864.
 60. " " " "
 61. Wash'n Ast. & Met. Obs'ns, 1865.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
50	0	From zenith to E.		
51		From the N. E.		
52		From zenith to N. W.		
53	Train visible 5s.....		From N. E. to N.		
54	Visible train.....		From E. to N. W.	Disappeared with an explosion.	
55	Visible train.....	45	From the S. E.	Seen also at Westtown and at Wilmington and Odessa, Del.	
56		From W. to E.		
57	Train 15' long.....		From the E. N. E.		
58	Train 3 or 4° degrees long.....		From Delphinus towards Pegasus.	Disappeared with a loud report.....	John Gardner.
59	Visible train.....	40	Obliquely towards horizon.		
60		Vertically to horizon.		
61	Very large and brilliant.	
62	Brilliant.	
63		Towards the N.		
64	A beautiful blue train.....	30	From E. to W.	Passed below some cirro-stratus clouds.....	David Trowbridge.
65	Report of explosion heard in 3/4 m.....	Isaac Coles.
66	
67	Brilliant.	J. K. Larimore.
68	Train 3 or 4° long.....		When first seen its altitude was 40°.....	Pliny Earle Chase.
69	25	Disappeared with an explosion.....	
70		Vertically to the horizon.....	Max. at appearance = η Tauri, at disappearance = Jupiter.	
71	Train visible 1m.....		Towards horizon at an angle of 75°.	Visible less than is. Disappeared with a report heard over an area 120 miles in diameter.	Prof. B. F. Mudge.
72		Vertically to horizon.	
73	Brilliant train.....	30	Burst into fragments with a loud explosion. Seen in many places in New England, New York, and New Jersey.	Described by Prof. E. Loomis.
74	Train visible 5m.		
75	Train 30 feet long, 3 feet wide.....		Seen by many observers.....	Described by Prof. J. L. Smith.

62. Wash'n Ast. & Met. Obs'ns, 1865.

63. " " " "

64. A. J. S. XLII, 286.

65. Smithsonian Archives (not yet printed).

66. Wash'n Ast. & Met. Obs'ns, 1867.

67. Wash'n Ast. & Met. Obs'ns, 1867.

68. A. J. S. XLIV, 288.

69. Proc. Am. Phil. Soc. X, 353.

70. Wash'n Ast. & Met. Obs'ns, 1868.

71. A. J. S. XLVI, 429.

72. Wash'n Ast. & Met. Obs'ns, 1869.

73. A. J. S. XLVIII, 145.

74. Wash'n Ast. & Met. Obs'ns, 1869, 372; Proc. Am. Phil. Soc. XI, 194.

75. A. J. S. XLIX, 139.

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day	Hour and minute.				
76	1869	Nov.	10	<i>h. m.</i> 6 53	Very bright	<i>s.</i> 4	In the N. W.
77	1869	Dec.	23	8 30	3d magnitude	White	In the E.
78	1870	Mar.	24	16 20	4th magnitude	White	0.4	
79	1870	Mar.	25	9 45	3d magnitude	White	0.3	
80	1870	Apr.	19	8 35	4th magnitude	White	2.4	
81	1870	Apr.	19	8 37	3d magnitude	White	0.5	
82	1870	Apr.	19	9 33	4th magnitude	White	0.5	
83	1870	Apr.	19	10 12	4th magnitude	White	0.5	
84	1870	Apr.	19	15 6	5th magnitude	White	0.5	
85	1870	Apr.	25	11 11	4th magnitude	White	0.5	
86	1876	Apr.	27	14 3	4th magnitude	White	0.4	
87	1870	June	19	10 12	5th magnitude	Reddish	0.3	
88	1870	June	21	10 0	2d magnitude	Reddish	2.0	
89	1870	June	21	10 0	6th magnitude	White	0.4	
90	1870	June	24	10 35	1st magnitude	Orange	1.5	
91	1870	June	25	9 23	2d magnitude	White	1.5	
92	1870	June	25	12 36	Size of Jupiter	Orange and green ..	2.5	Midway between Spica and Antares.
93	1870	June	25	12 36	6th magnitude	White	0.3	
94	1870	July	26	9 50	Size of Jupiter	White	0.2	
95	1870	July	30	8 50	3d magnitude	White	3.0	From 240° + 75° to 195° + 75°.
96	1870	July	30	9 14	Size of Saturn	White	0.2	From 14h. 50m. + 20°.
97	1870	July	30	9 20	3d magnitude	White	3.0	From 17h. 0m. — 25°.
98	1870	July	30	9 40	4th magnitude	White	1.5	From 4° south of Saturn.
99	1870	July	30	9 50	2d magnitude	White	0.2	From a Aquilæ.
100	1870	Aug.	15	9 54	4th magnitude	White	0.5	From a Aquarii.
101	1870	Sept.	4	8 5	4th magnitude	Orange	0.5	From a Pegasi.
102	1870	Sept.	24	8 15	5th magnitude	White	0.4	From 10° N. of Arcturus.
103	1870	Sept.	24	8 30	3d magnitude	White	0.4	
104	1870	Oct.	15	8 10	3d magnitude	White	0.3	From 19h. 30m. + 60° directly tow' ds Polaris.
76.	Wash'n Ast. & Met. Obs'ns, 1869.							
77.	Wash'n Ast. & Met. Obs'ns, 1870.							
78.	Wash'n Ast. & Met. Obs'ns, 1870.							
79.	Wash'n Ast. & Met. Obs'ns, 1870.							
80.	Wash'n Ast. & Met. Obs'ns, 1870.							
81.	Wash'n Ast. & Met. Obs'ns, 1870.							
82.	Wash'n Ast. & Met. Obs'ns, 1870.							
83.	Wash'n Ast. & Met. Obs'ns, 1870.							
84.	Wash'n Ast. & Met. Obs'ns, 1870.							
85.	Wash'n Ast. & Met. Obs'ns, 1870.							
86.	Wash'n Ast. & Met. Obs'ns, 1870.							
87.	Wash'n Ast. & Met. Obs'ns, 1870.							
88.	Wash'n Ast. & Met. Obs'ns, 1870.							
89.	Wash'n Ast. & Met. Obs'ns, 1870.							
90.	Wash'n Ast. & Met. Obs'ns, 1870.							

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
76	No train.....	0	N. 20° E.....	From δ Aquilæ.....	Prof. J. R. Eastman.
77	No train.....	8	S. to N.....	Across the "Sickle".....	Prof. J. R. Eastman.
78	No train.....	5	S. 5° W.....	From a point on the meridian 10° S.....	Prof. J. R. Eastman.
80	Short train.....	10	S. W.....	South of α Canis Majoris.....	Prof. J. R. Eastman.
81	No train.....	4	Northerly from near the eastern horizon.....	N. Cahill.
82	No train.....	Northerly from near the eastern horizon.....	N. Cahill.
83	No train.....	Northerly from near the eastern horizon.....	N. Cahill.
84	No train.....	10	Towards the W.....	Across the "Dipper".....	N. Cahill.
85	No train.....	10	Towards the E.....	From 10° south of the zenith.....	D. Horgan.
86	No train.....	8	N. W. from α Cygni.....	D. Horgan.
87	No train.....	12	N. from α Bootis.....	D. Horgan.
88	Long train, lasting 3s.....	D. Horgan.
89	No train.....	5	South.....	From 5 Ursæ Minoris to 8° south of α Bootis.....	D. Horgan.
90	No train.....	10	Easterly.....	From alt. 40° through the constellation of the Scorpion.....	Prof. J. R. Eastman.
91	No train.....	8	West.....	Prof. J. R. Eastman.
92	Train visible 0.8s.....	30	West.....	From β Cassiopeiæ to 10° below Polaris.....	Prof. J. R. Eastman.
93	No train.....	4	East.....	Exactly across Polaris in a line parallel to the above course.....	Prof. J. R. Eastman.
94	No train.....	45	West.....	Prof. J. R. Eastman.
95	No train.....	6	West.....	Prof. J. R. Eastman.
96	No train.....	15	West.....	Prof. J. R. Eastman.
97	No train.....	15	West.....	Between 9h. and 10h. saw ten other small meteors, all moving westward.	Prof. J. R. Eastman.
98	No train.....	6	South.....	Prof. J. R. Eastman.
99	No train.....	8	West.....	Prof. J. R. Eastman.
100	Train visible 0.6s.....	10	South west.....	D. Horgan.
101	No train.....	8	Vertically to horizon.....	D. Horgan.
102	No train.....	8	East.....	"Across the Dipper".....	D. Horgan.
103	Train visible 0.3s.....	10	"From Polaris towards the Dipper".....	D. Horgan.
104	No train.....	4	Prof. J. R. Eastman.
91. Wash'n Ast. & Met. Obs'ns, 1870.					
92.	"	"	
93.	"	"	
94.	"	"	
95.	"	"	
96. Wash'n Ast. & Met. Obs'ns, 1870.					
97.	"	"	
98.	"	"	
99.	"	"	
100.	"	"	
101. Wash'n Ast. & Met. Obs'ns, 1870.					
102.	"	"	
103.	"	"	
104.	"	"	

CATALOGUE V.-Sporadic Meteors-Continued.

Ret. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day.					
105	1870	Nov. ...	3	h. m.	3d magnitude	White	8.	From α Andromedæ.
106	1870	Nov. ...	14	11 29	3d magnitude	White	0.3	From Orion towards Sirius.
107	1870	Nov. ...	15	11 50	1st magnitude	White	0.5	
108	1870	Nov. ...	15	7 18	3d magnitude	White	2.0	
109	1870	Nov. ...	15	7 25	3d magnitude	White	0.5	
110	1870	Dec. ...	13	9 5	3d magnitude	White	2.0	
111	1870	Dec. ...	13	9 30	3d magnitude	White	0.2	
112	1870	Dec. ...	13	12 15	3d magnitude	White		
113	1870	Dec. ...	13	12 36	1st magnitude	Orange	3.0	From 8° N. of Jupiter.
114	1871	May ...	23	9 43	1st magnitude	White	3.5	On the meridian, Z. D. S. 55° .
115	1871	July ...	6	8 35	2d magnitude	White	1.0	8° N. of Antares.
116	1871	July ...	6	8 38	2d magnitude	White	1.5	
117	1871	July ...	6	8 45	4th magnitude	White	0.8	15° N. W. of Antares.
118	1871	July ...	6	8 55	1st magnitude	White	3.0	
119	1871	Aug. ...	19	8 or 9	" Large ball of fire "	Crimson.		
120	1871	Aug. ...	8	11 55	2d magnitude	White	1.0	40° N. of Algol.
121	1871	Aug. ...	9	10 45	2d magnitude	White	0.3	Near β Pegasi.
122	1871	Aug. ...	12	11 53	1st magnitude	White and orange	1.0	At α Aquilæ.
123	1871	Aug. ...	16	8 26	As large as Sirius			From Aldebaran passed 5° below the Pleiades.
124	1871	Aug. ...	16	8 48	Larger than Jupiter			
125	1871	Nov. ...	16	13 55	1st magnitude	White		
126	1872	Jan. ...	22	8 15	1st magnitude	Red, then white	10m.	
127	1872	July ...	30	9 5	1st magnitude	Orange	4	From β Cassiopeiæ towards Polaris.
128	1872	July ...	28	8 55	1st magnitude	White	1	
129	1872	Aug. ...	28	9 18	1st magnitude	White	1	
130	1872	Aug. ...	3	10 20	2d magnitude	White	1	20° E. of Polaris.
131	1872	Aug. ...	6	9 55	2d magnitude	White	1	At Polaris.
132	1872	Aug. ...	9	8 18	2d magnitude	Orange and white	2	
133	1872	Aug. ...	9	8 37	2d magnitude	White	1.5	
134	1872	Aug. ...	9	8 40	3d magnitude	White	1	
105.	Wash'n Ast. & Met. Obs'ns, 1870.				110. Wash'n Ast. & Met. Obs'ns, 1870.			115. Wash'n Ast. & Met. Obs'ns, 1871.
106.	" "				111. " "			116. " "
107.	" "				112. " "			117. " "
108.	" "				113. Wash'n Ast. & Met. Obs'ns, 1871.			A. J. S. II, 227.
109.	" "				114. " "			118. Wash'n Ast. & Met. Obs'ns, 1871.
								119. Wash'n Ast. & Met. Obs'ns, 1871.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
105	Train visible 0.4s.....	0	Southwest.....	D. Horgan.
106	Train visible 0.3s.....	5	D. Horgan.
107	Train visible 2.0s.....	8	West.....	Passed south of α Pegasi.....	D. Horgan.
108	No train.....	25	South from α Cygni.....	D. Horgan.
109	No train.....	8	I. Horgan.
110	Train visible 0.3s.....	4	N. Cahill.
111	22	N. Cahill.
112	Short train.....	8	Westerly.....	N. Cahill.
113	15	Westerly.....	N. Cahill.
114	15	Westerly.....	Prof. J. R. Eastman.
115	18	West from Antares.....	Prof. J. R. Eastman.
116	10	South.....	Prof. J. R. Eastman.
117	35	West from ϵ Cassiopeæ.....	Prof. J. R. Eastman.
118	12	From South to North.....	Very brilliant.....	Prof. J. R. Eastman.
119	Short train, visible 7s.....	12	From α to β Aurigæ.....	Burst into three fragments with a loud explosion.....	E. S. Martin.
120	8	South.....	D. Horgan.
121	Train visible 2s.....	15	South.....	Very brilliant.....	Prof. J. R. Eastman.
122	20	South.....	D. Horgan.
123	Train visible 1s.....	35	South.....	Very brilliant.....	Prof. J. R. Eastman.
124	Train visible 3s.....	25	South.....	Prof. J. R. Eastman.
125	20	From West to East.....	Seen at several other cities in Mexico.....	Prof. A. Hall.
126	8	Fell vertically from α Cassiopeæ.....	Motion seemed to be by impulses, but slow.....	José de la Vega.
127	Short train.....	12	South.....	D. Horgan.
128	15	Towards γ Ursa Majoris.....	D. Horgan.
129	Short train.....	8	N. E. from Saturn.....	D. Horgan.
130	Long train.....	15	S. from α Cassiopeæ.....	D. Horgan.
131	20	S. from ζ Ursa Majoris.....	D. Horgan.
132	Long train.....	10	D. Horgan.
133	Short train.....	10	D. Horgan.

130. Wash'n Ast. & Met. Obs'ns, 1872.

131. " " " " " "

132. " " " " " "

133. " " " " " "

125. A. J. S. III, 235.

126. Wash'n Ast. & Met. Obs'ns, 1872.

127. " " " " " "

128. " " " " " "

129. " " " " " "

120. Wash'n Ast. & Met. Obs'ns, 1871.

121. " " " " " "

122. " " " " " "

123. " " " " " "

124. " " " " " "

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Hour and minute.					
134	1872	Aug. ...	<i>h. m.</i>	U. S. Naval Observatory...	2d magnitude	White	8.	Appeared in the cluster in sword handle of Perseus.
135	1872	Aug. ...	9 5	U. S. Naval Observatory...	2d magnitude	Orange	1	
136	1872	Aug. ...	9 16	U. S. Naval Observatory...	2d magnitude	Yellow-red	1	
137	1872	Nov. ...	9 58	U. S. Naval Observatory...	2d magnitude	Orange	1.5	From 9° E. of α Orionis vertically towards the horizon
138	1872	Dec. ...	9 0	U. S. Naval Observatory...	1st magnitude	White	1.0	From ζ Ursæ Minoris towards ζ Draconis.
139	1872	Dec. ...	10 45	Bloomington, Ind.	"Large"	Red	Altitude 40°, 15° E. of meridian.
140	1872	Dec. ...	12 53	Louisville, Ky.	1st magnitude	White	Appeared in the zenith.
141	1873	Jan. ...	3 0	U. S. Naval Observatory...	One green; the following yellow.	1	Appeared at γ Orionis.
142	1873	Feb. ...	14 6 0	New Haven, Conn.	White	One green; the following yellow.
143	1873	April ...	3 11 5	U. S. Naval Observatory...	1st magnitude	White	1.5	Appeared near α Virginis.
144	1873	May ...	8 5	Harbor Grace, Newfoundland.	W. S. W. at an altitude of 40°.
145	1873	June ...	6 9 36	U. S. Naval Observatory...	1st magnitude	Orange	1.5	From Polaris vertically towards horizon.
146	1873	June ...	6 9 37	U. S. Naval Observatory...	1st magnitude	Orange	2.0	From Polaris 8° towards δ Ursæ Majoris.
147	1873	Aug. ...	12 17	U. S. Naval Observatory...	1st magnitude	Orange	1 5	From ζ Ursæ Minoris south at an angle of 30° with the horizon.
148	1873	Aug. ...	10 8 10	U. S. Naval Observatory...	1st magnitude	Orange	3.0	Appeared at θ Cygni.
149	1873	Nov. ...	6 0	U. S. Naval Observatory...	Bright as new moon	Intense white	2.0	Appeared at θ Aquilæ.
150	1873	Dec. ...	14 7 30	U. S. Naval Observatory...	2d magnitude	White	4.0	From 12° E. of Polaris vertically towards the horizon.
151	1873	Dec. ...	7 39	Washington, D. C.	Entered the atmosphere over the State of Delaware at a height of 90 miles, and disappeared over Fairfax Co., Va., at a height of from 8 to 20 miles.
152	1874	May ...	9 13	U. S. Naval Observatory...	1st magnitude	Orange and violet	3	Appeared near Polaris.
153	1874	July ...	18 10 0	Louisville, Ky.	One-third diam. of moon.	Greenish white	3	Appeared 5° above Mars and passed 8° W. of Mars.
154	1874	Aug. ...	10 15 20	U. S. Naval Observatory...	15' in diameter	White and orange	3.5	
155	1875	July ...	9 45	U. S. Naval Observatory...	Bright as Mars	

134. Wash'n Ast. & Met. Obs'ns, 1872.

135. " " " " " "

136. " " " " " "

137. " " " " " "

138. Wash'n Ast. & Met. Obs'ns, 1872.

139. A. J. S. V., 318.

140. A. J. S. V., 203.

141. Wash'n Ast. & Met. Obs'ns, 1873.

142. A. J. S. V., 318.

143. Wash'n Ast. & Met. Obs'ns, 1873.

144. A. J. S. V., 154.

145. Wash'n Ast. & Met. Obs'ns, 1873.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
134	Short train	9	S. from ϵ Ursa Majoris	D. Horigan.
135	15	S. from α Andromedæ	D. Horigan.
136	Orange train	To γ E. of Polaris	Prof. J. R. Eastman.
137	15	Prof. J. R. Eastman.
138	10	Prof. J. R. Eastman.
139	E. of S., making an angle of 60° with the horizon.	Seen also at several places in Kentucky	Prof. D. Kirkwood.
140	Left a dense stream of blue smoke, which remained visible several minutes.	Towards the horizon in a southerly direction.	Seen from several other points, and at some places its disappearance was followed by loud detonations. (Probably same meteor as preceding.)	Prof. J. L. Smith.
141	Short train	10	Vertically towards the horizon. North and down to an alt. of 15° in the direction N. 64° W.	D. Horigan.
142	Vertically towards the horizon. To W. N. W. at altitude of 10° ..	Two meteors close together. Seen also at New Britain, Conn.	Prof. H. A. Newton.
143	Yellow train	10	D. Horigan.
144	8	Burst, leaving a coppery-red cloud visible 30m.	Henry H. Cleft.
145	Prof. J. R. Eastman.
146	15	Prof. J. R. Eastman.
147	Short train	Prof. J. R. Eastman.
148	Short train	45	Across α Aquilæ	Prof. J. R. Eastman.
149	Train separated into two distinct parts.	A. N. Skinner.
150	No train	15	Prof. J. R. Eastman.
151	This meteor was seen by many observers from Delaware to Virginia, and all the available data were collected and discussed by Cleveland Abbe in a report to the Washington Philosophical Society.	
152	Long train	30	Southeasterly	When within 20° of the horizon it burst into three or four fragments, flashing forth red and blue lights.	D. Horigan.
153	From N. W. to S. E.	Seen through the slit of the great dome from zenith towards S. E.	Prof. J. L. Smith.
154	Continuous train	30	Prof. E. S. Holden.
155	Continuous train	15	Fell vertically	Separated into two parts, each of which moved 5° before extinction.	D. Horigan.
146.	Wash'n Ast. & Met. Obs'ns, 1873.	150. Wash'n Ast. & Met. Obs'ns, 1873.	
147.	" " " "	"	"	151. Bull. Wash. Phil. Soc. II, 139.	
148.	" " " "	"	"	152. Wash'n Ast. & Met. Obs'ns, 1874.	
149.	" " " "	"	"	153. A. J. S. X, 203. 154. Wash'n Ast. & Met. Obs'ns, 1874. 155. Wash'n Ast. & Met. Obs'ns, 1875.	

CATALOGUE V.-Sporadic Meteors-Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day, hour and minute.					
156	1875	July ...	14	U. S. Naval Observatory ...	Large as Mars.....	Orange and red	s. 4	Appeared 10° above Mars and passed 5° E. of Mars.
157	1875	Aug. ...	5	U. S. Naval Observatory ...	2d magnitude	Orange	2	Moved from λ Sagittarii towards λ Scorpii.
158	1875	Sept. ...	16	U. S. Naval Observatory ...	Three times size of Venus.	White	3	Moved from Saturn to λ Aquarii.
159	1875	Oct. ...	17	U. S. Naval Observatory ...	1st magnitude ...	White and green ...	2	Appeared near β Aquarii.
160	1875	Nov. ...	14	U. S. Naval Observatory ...	Size of Mars	White	2	Appeared in Cassiopeæ.
161	1875	Nov. ...	17	U. S. Naval Observatory ...	3d magnitude	White	1	Appeared near Polaris.
162	1875	Nov. ...	7	U. S. Naval Observatory ...	1st magnitude	White	0.7	Appeared at ξ Ceti.
163	1875	Nov. ...	7	U. S. Naval Observatory ...	1st magnitude	White	0.7	Appeared at χ Ceti.
164	1875	Nov. ...	17	U. S. Naval Observatory ...	1st magnitude	White	0.7	Appeared at χ Ceti.
165	1875	Nov. ...	22	U. S. Naval Observatory ...	8' to 12' in diameter	Violet or pink in front, bright green in center, and pale green following.	6 to 8	Appeared at χ Ceti.
166	1875	Dec. ...	27	Few sec's.	From W. to E. across the northern sky.
167	1876	Jan. ...	26	U. S. Naval Observatory ...	2d magnitude	White	0.7	From γ Canis Majoris to ϵ Canis Majoris.
168	1876	Jan. ...	31	Louisville, Ky.	$\frac{1}{2}$ diameter of moon.....	2 or 3	60° above the N. W. horizon.
169	1876	April ...	19	U. S. Naval Observatory ...	3d magnitude	Orange and red	1
170	1876	July ...	8	Appeared at an altitude of 88 miles.
171	1876	Aug. ...	8	U. S. Naval Observatory ...	2d magnitude	White and green ...	2	S. W. from ϵ Urs. Min. towards Arcturus.
172	1876	Aug. ...	9	U. S. Naval Observatory ..	2d magnitude	White and green ...	3	S. W. from midway between Polaris and Cassiopeæ.
173	1876	Aug. ...	13	U. S. Naval Observatory ...	3d magnitude	White	1	Appeared 8° below Polaris.
174	1876	Aug. ...	14	U. S. Naval Observatory ...	3d magnitude	White	1
175	1876	Aug. ...	15	U. S. Naval Observatory ...	3d magnitude	White	1
176	1876	Aug. ...	15	U. S. Naval Observatory ...	3d magnitude	White	1
177	1876	Aug. ...	15	U. S. Naval Observatory ...	3d magnitude	White	1
178	1876	Aug. ...	21	U. S. Naval Observatory ...	4th magnitude	White	0.5
179	1876	Aug. ...	22	U. S. Naval Observatory ...	3d magnitude	White	1
180	1876	Aug. ...	27	U. S. Naval Observatory ...	3d magnitude	White	1	Appeared 5° below α Cassiopeæ.

156. Wash'n Ast. & Met. Obs'ns, 1875.

157. " " "

158. " " "

159. Wash'n Ast. & Met. Obs'ns, 1875.

160. " " "

161. " " "

162. Wash'n Ast. & Met. Obs'ns, 1875.

163. " " "

164. " " "

CATALOGUE V.-Sporadic Meteors-Continued.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
156	Continuous train.....	0			D. Horigan.
157	Slight train.....	12	Fell vertically.....	Exploded and illuminated the small cloud behind which it passed.	Prof. J. R. Eastman.
158	Slight train.....	15			D. P. Todd.
159	Train visible 20s.....	18			
160	No train.....	25	Fell vertically.....		D. P. Todd.
161	Short train.....	32	Fell nearly vertically.....		D. Horigan.
162	No train.....	10	Fell vertically.....		D. P. Todd.
163	No train.....		Fell vertically.....	All disappeared at an altitude of 5°.....	D. P. Todd.
164	No train.....		Fell vertically.....		D. P. Todd.
165	Pale greenish train.....		Moved vertically through Cygni and nearly through β Cygni.	Within 6° or 8° of the horizon it passed behind a bank of clouds.	Prof. E. S. Holden.
166	Faint train.....	8		Seen in Missouri, Kansas, Iowa, Nebraska, and Ohio.....	G. C. Brodhead.
167	Train short, green.....		From N. W. to S. E.....		D. P. Todd.
168	Train visible 40m.....	15	West from Jupiter.....	Seen by several other observers in Kentucky.....	Prof. J. L. Smith.
169			From N. 78° W. over N. E. corner of Indiana.		D. Horigan.
170				Exploded over Lake Michigan at an altitude of 34 miles.....	Prof. D. Kirkwood.
171	Long train.....	20			D. Horigan.
172	Train lasted 13s.....	30	Westward.....		D. Horigan.
173	Long train, visible 2s.....	15	South from α Andromedæ.....		D. Horigan.
174	Short train.....	20	"South from the Dipper".....		G. Anderson.
175	Short train.....	15	Eastward.....		D. Horigan.
176	Short train.....	15	West across Polaris.....		D. Horigan.
177	Long train.....	20	N. E. from β Pegasi.....		D. Horigan.
178	Short train.....	15	S. W. from Polaris.....		D. Horigan.
179	Short train.....	12	E. S. E. from Saturn.....		D. Horigan.
180	Short train.....	10			D. Horigan.

165. Wash'n Ast. & Met. Obs'ns, 1875.
 166. Trans. St. Louis Acad. Sci. III, 349.
 167. Astron. and Meteor. Observations, 1876.
 168. A. J. S. XIV, 468.
 169. Wash'n Ast. & Met. Obs'ns, 1876.
 170. A. J. S. XIV, 75.
 171. Wash'n Ast. & Met. Obs'ns, 1876.
 172. " " " "
 173. " " " "
 174. " " " "
 175. " " " "
 176. Wash'n Ast. & Met. Obs'ns, 1876.
 177. " " " "
 178. " " " "
 179. " " " "
 180. " " " "

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
181	Short train	0	West from Polaris.....	D. Horgan.
182	Train of short duration	10 or 15	From N. towards N. E.....	Prof. E. S. Holden.
183	8	Fell vertically to horizon.....	D. Horgan.
184	8	Fell vertically to horizon.....	D. Horgan.
185	Train 1.5° long, lasted 1s.....	15	Down and north.....	Like the preceding, but 10° further north.....	D. P. Todd.
186	Followed by a great number of small meteors.	From N. W. to N. E.....	Passed within 30' of moon's south limb; moon nearly full. Seen in Kansas, Missouri, Illinois, Indiana, and Ohio; disappeared with an explosion.	Prof. D. Kirkwood.
187	Train visible several seconds	J. S. Hunter.
188	4	Westerly	3d magnitude at appearance and three times magnitude of Sirius at disappearance.	D. P. Todd.
189	15	D. P. Todd.
190	Light train.....	30	Eastward.....	Disappeared at altitude of 12° or 13°.....	Prof. J. R. Eastman.
191	20	Vertically downwards.....	F. M. Parker.
192	Observer thought he heard the sound of the explosion.....	Prof. E. S. Holden.
193	10 to 12	Towards the west at an angle of 65° with the vertical.....	A. N. Skinner.
194	12 to 15	Fell nearly vertical.....	Robert C. Hindley.
195	To 4° above β Cassiopee.....	Probably same as preceding.....	Dr. F. L. James.
196	A bright train.....	11+	Near α Cassiopee it broke into seven or eight fragments.....	E. Colbert.
197	6	Across α Cygn.....	G. W. E. Tronvelot.
198	10	Rapid	E. F. Sawyer.
199	5	Quite slow.....	E. F. Sawyer.
200	21	Rapid	E. F. Sawyer.
201	21	Slow	E. F. Sawyer.
202	21	Across ϵ and ζ Aquilæ	Rapid, probably a Perseid	E. F. Sawyer.
203	21	E. F. Sawyer.
204	Red streak	23+	Rapid	E. F. Sawyer.
205	Quite slow.....	E. F. Sawyer.
187.	Proc. Am. Phil. Soc. XVI, 566.	194. Proc. Am. Phil. Soc. XVII, 340.	200. A. J. S. XVI, 348.	
188.	Wash'n Ast. & Met. Obs'ns, 1877.	195. " " " "	201. " " " "	
189.	" " " "	196. Proc. Am. Phil. Soc. XVIII, 239.	202. " " " "	
190.	" " " "	197. Proc. Am. Phil. Soc. XVIII, 239.	203. " " " "	
191.	A. J. S. XIV, 163.	198. A. J. S. XVI, 348.	204. " " " "	
192.	Wash'n Ast. & Met. Obs'ns, 1877.	199. " " " "	205. " " " "	
193.	" " " "	

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.	
	Year.	Month.	Day.						
			h. m.						
206	1878	Aug.	3	9 49	Cambridge, Mass.	1st magnitude	8.	From 340°	24° to 344° + 33°.
207	1878	Aug.	3	11 22	Cambridge, Mass.	1st magnitude	From 19	+ 30 to 22 + 47.
208	1878	Aug.	3	11 27	Cambridge, Mass.	1st magnitude	From 47	+ 55 to 50 + 50.
209	1878	Aug.	11	10 10	Bloomington, Ind.	One-third size of moon	2	Appeared in the E. at an altitude of 10°.	
210	1878	Aug.	20	8 32	Cambridge, Mass.	1st magnitude	From 155°	+ 72½° to 137½° + 70°.
211	1878	Aug.	22	8 59	Cambridge, Mass.	1st magnitude	From 185	+ 70 to 170 + 68.
212	1878	Aug.	22	10 2	Cambridge, Mass.	Equal to Mars.	3	From 185	+ 63 to 230 + 77.
213	1878	Aug.	25	8 59	Cambridge, Mass.	1st magnitude	From 249½	+ 39 to 250½ + 27.
214	1878	Aug.	27	10 4	Cambridge, Mass.	1st magnitude	From 263½	+ 48 to 245 + 59½.
215	1878	Aug.	27	10 19	Cambridge, Mass.	1st magnitude	From 245	+ 56 to 229 + 58.
216	1878	Aug.	29	8 46	Cambridge, Mass.	1st magnitude	From 187½	+ 45 to 185½ + 38½.
217	1878	Aug.	30	8 32	Cambridge, Mass.	1st magnitude	From 297	+ 50 to 252½ + 62½.
218	1878	Aug.	31	8 35	Cambridge, Mass.	1st magnitude	From 212	+ 24 to 202½ + 26.
219	1878	Sept.	1	10 8	Cambridge, Mass.	1st magnitude	From 236	+ 30 to 232 + 22.
220	1878	Sept.	16	9 0	Henryville, Clark Co., Ind.	One-fourth or one-fifth diameter of moon.	From γ Ursæ Majoris over δ Ursæ Majoris.	
221	1878	Sept.	18	8 29	Cambridge, Mass.	1st magnitude	From 323°	15° to 325° - 20°.
222	1878	Sept.	20	9 39	Cambridge, Mass.	1st magnitude	From 329	- 2 to 326 - 16.
223	1-78	Sept.	22	7 19	Cambridge, Mass.	1st magnitude	From 289	- 15 to 295 - 2.
224	1878	Sept.	22	8 33	Cambridge, Mass.	Larger than Jupiter.	2	From 2½	- 17½ to 5 - 22.
225	1878	Sept.	23	7 18	Cambridge, Mass.	1st magnitude	From 355½	+ 12½ to 346 - 6½.
226	1878	Sept.	23	8 25	Cambridge, Mass.	1st magnitude	From 30	+ 28 to 10 + 26.
227	1878	Sept.	23	9 26	Cambridge, Mass.	Larger than 1st mag.	2	From 42	+ 56 to 105 + 65.
228	1878	Sept.	27	8 10	Cambridge, Mass.	1st magnitude	1.5	From 41½	+ 16 to 347 + 23.
229	1878	Sept.	28	9 52	Cambridge, Mass.	1st magnitude	From 49	+ 18 to 47 - 2.
230	1878	Sept.	29	9 5	Cambridge, Mass.	1st magnitude	From 15	+ 3 to - - stationary.
231	1878	Sept.	29	9 36	Cambridge, Mass.	1st magnitude	From 48	+ 21 to 55 + 17½.
232	1878	Oct.	28	6 44	Boston, Mass.	Nearly as bright as Mars.	1	From 31½	+ 37 to 20 + 37.
233	1878	Nov.	12	7 0	Washington, Ind.	Two-thirds diam. of moon.	10	From a Lyrae to 20° N. W. of Jupiter.	
234	1878	Nov.	12	7 15	Boston, Mass.	As brilliant as Sirius.	Through the trapezium in Ursæ Minor.	
235	1878	Nov.	13	6 40	New Haven, Conn.	1st or 2d magnitude.	1 to 1.5	From a Lyrae to near χ Ophiuchi.	

206. A. J. S. XVI₃, 348.

207. " " "

208. A. J. S. XVI₃, 348.

209. Proc. Am. Phil. Soc. XVIII, 240.

210 to 219. A. J. S. XVI₃, 348.

220. Proc. Am. Phil. Soc. XVIII, 241.

221 to 231. A. J. S. XVI₃, 348.

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	DATE.			Place of observation.	Apparent size.	Color.	Duration.	Position, altitude and azimuth.
	Year.	Month.	Day.	Hour and minute.				
236	1878	Nov. ...	14	<i>h. m.</i> 3 30	<i>s.</i>	South of observer. From the Crab Nebula to above and to the left of α Ursa Majoris. S. 10° W.; alt., 25° .
237	1878	Dec. ...	19	9 30	Very bright	3	
238	1878	Dec. ...	30	7	Brilliant.	
239	1879	Jan. ...	27	14 28	Larger than moon.	
240	1879	Mar. ...	14	15 53	Pale blue.....	

CATALOGUE V.—Sporadic Meteors—Continued.

Ref. number.	Train, appearance and duration.	Length of path.	Direction of motion.	Remarks.	Observer.
236	Seen during bright sunshine	Thomas Whitaker.
237	J. J. Skinner.
238	Seen in Indiana, Pennsylvania, and Ohio	Prof. D. Kirkwood.
239	Explosion heard 4m. after disappearance.....	Thomas T. Bates.
240	Train visible several minutes.	From S. W. to N. E.....	Disappeared with an explosion	Prof. D. Kirkwood.
236. Proc. Am. Phil. Soc. XVIII, 241.					
237. Science Observer, II, 35.					
238. Proc. Am. Phil. Soc. XVIII, 241.					
239. Proc. Am. Phil. Soc. XVIII, 243.					
240. Proc. Am. Phil. Soc. XVIII, 245.					





14 DAY USE
RETURN TO DESK FROM WHICH BORROWED

LOAN DEPT.

This book is due on the last date stamped below, or
on the date to which renewed.
Renewed books are subject to immediate recall.

9 May '62 G N

DEC 8 1967 77

RECEIVED

REC'D LD NOV 26 '67 -2 PM

MAY 13 1962 LOAN DEPT.

ASTRON., MATH,
STAT. LIBRARY

DEC 10 1969

NOV 9 1967 77

RECEIVED

NOV 13 '67 -12 M

LOAN DEPT.

DEC 9 1969 5 6

REC'D LD DEC 11 '69 -7 PM

LD 21A-50m-3, '62
(C7097s10) 476H

General Library
University of California
Berkeley

M512210

QB741

Eastman, J.R.

E2

The progress of
meteoric astronomy in
America

M512210

QB741
E2

